

WEAK NORMALITY OF FAMILIES OF MEROMORPHIC MAPPINGS AND BUBBLING IN HIGHER DIMENSIONS

S. IVASHKOVICH, F. NEJI

ABSTRACT. Our primary goal in this paper is to understand whether the sets of normality of families of meromorphic mappings between general complex manifolds are pseudoconvex or not. It turns out that the answer crucially depends on the type of convergence one is interested in. We examine three natural types of convergence introduced by one of us earlier and prove pseudoconvexity of sets of normality for a large class of target manifolds for the so called *weak convergence*. Furthermore we determine the structure of the exceptional components of the limit of a weakly but not strongly converging sequence - they turn to be *rationally connected*. This observation allows to determine effectively when a weakly converging sequence fails to converge strongly. An application to the Fatou sets of meromorphic self-maps of compact complex surfaces is given.

CONTENTS

1. Introduction	1
2. Topologies on the space of meromorphic mappings	6
3. Pseudoconvexity of sets of normality	10
4. Convergence of mappings with values in projective space	13
5. Bloch-Montel type normality criterion	17
6. Behavior of volumes of graphs under weak and gamma convergence	21
7. Rational connectivity of the exceptional components of the limit	23
8. Fatou components	26
References	31

1. INTRODUCTION

1.1. Convergence of meromorphic mappings. Domains of convergence of meromorphic functions where, probably, for the first time considered by Cartan and Thullen in [CT] simultaneously with their domains of existence. In particular in [CT] it was proved that these domains are pseudoconvex. Later domains of convergence of meromorphic functions were studied in [Ru]. Already from these sources it is clear that when working with sequences of meromorphic functions and, more generally, mappings one finds himself bounded to consider several notions of their convergence. Some of these notions were introduced in [Fu], [Iv2], we shall recall the essentials in section 2. It occurs that pseudoconvexity or not of domains of convergence/normality in the case of meromorphic

Date: January 20, 2013.

1991 *Mathematics Subject Classification.* Primary - 32H04, Secondary - 32H50, 32Q45.

Key words and phrases. Meromorphic mapping, convergence, normal family.

mappings crucially depends on the type of convergence one is looking for. Let briefly describe the ways one can define what does it means that a sequence $\{f_k\}$ of meromorphic mappings between complex manifolds U and X converge.

We start with the most obvious one. A sequence $\{f_k\}$ of meromorphic mappings between complex manifolds U and X is said to converge **strongly** to a meromorphic map f if the graphs Γ_{f_k} converge over compacts in U to the graph Γ_f as analytic sets. Convergence is meant here in Hausdorff metric, but as our first result shows they converge in a stronger topology of cycles.

Theorem 1. *If f_k strongly converge to f then for every compact $K \Subset U$ the volumes $\Gamma_{f_k} \cap (K \times X)$ are uniformly bounded and therefore Γ_{f_k} converge to Γ_f in the topology of cycles.*

This type of convergence is natural and has some nice features. For example the strong limit of a sequence of holomorphic maps is holomorphic and vice versa, if the limit f is holomorphic then for every compact $K \Subset U$ all f_k for $k \gg 1$ are holomorphic in a neighborhood of K and uniformly converge there to f . This statement was called the Rouché Principle in [Iv2].

But strong convergence has also some disadvantages. The first, crucial for us is the fact that the sets of strong convergence, i.e., maximal open subsets of U where a given sequence converge strongly on compacts, are not pseudoconvex in general. Moreover, the sets of strong normality (see later on) of families of meromorphic mappings can be just arbitrary, see Example 2.1. Also if one takes $X = \mathbb{P}^N$ the "most immediate" notion of convergence doesn't correspond to the strong one.

Therefore in [Iv2] along with the notion of strong convergence we proposed two weaker ones. We say that f_k converge **weakly** to f if they converge strongly to f on compacts outside of some analytic set A in U of codimension at least two. It turns out that this A can be taken to be the set I_f of points of indeterminacy of the limit map f and then for every compact K in $U \setminus I_f$ all weakly converging to f mappings f_k will be holomorphic on K (for k big enough) and converge to f uniformly on K , see Remark 3.1.

One more notion of convergence from [Iv2], which we need to recall here, is the gamma convergence (Γ -convergence). We say that f_k **gamma**-converge to f if they strongly converge to f outside of an analytic set (now it can be of codimension one) and for every divisor H in X and every compact $K \Subset U$ the intersections $f_k^* H \cap K$ have bounded volume counted with multiplicities, see more about the last condition in section 3.2.

Remark 1. Strong convergence (or **s**-convergence) will be denoted by $f_k \rightarrow f$, a weak one (or **w**-convergence) as $f_k \rightharpoonup f$, and Γ -convergence as $f_k \xrightarrow{\Gamma} f$. Note that in the second and third definitions we suppose that the limit f is defined and meromorphic on the whole of U if, even, the convergence takes place only on some part of U . In the first case the limit exists on the whole of U automatically, see Example 2.1 in concern with this.

For the better understanding of these notions let us give a description of the listed types of convergence in the case when X is projective, i.e., imbeds into \mathbb{P}^N for some N . In that special case the notions of convergence listed above permit an explicit analytic description as follows. Every meromorphic mapping f with values in \mathbb{P}^N can be locally represented as

$$f(z) = [f^0(z) : \dots : f^N(z)] \quad (1.1)$$

with holomorphic f^0, \dots, f^N , see section 4. More precisely, if $f : U \rightarrow \mathbb{P}^N$ is a meromorphic mapping then for every point $x_0 \in U$ there exists a neighborhood $V \ni x_0$ and holomorphic functions f^0, \dots, f^N in V satisfying (1.1). If the zero sets of f^j contain a common divisor then we can divide all f^j by its equation and get a representation such that $GCD(f^1, \dots, f^N) = 1$ in every \mathcal{O}_x , $x \in V$. In that case the indeterminacy set of f is

$$I_f \cap V = \{z \in V : f^0(z) = \dots = f^N(z) = 0\} \quad (1.2)$$

and has codimension at least two. Representation (1.1) satisfying (1.2) we shall call *reduced*.

We shall prove the following

Theorem 2. *Let $\{f_k\}$ be a sequence of meromorphic mappings from a complex manifold U to \mathbb{P}^N . Then:*

- i) $f_k \xrightarrow{\Gamma} f$ if and only if for any point $x_0 \in U$ there exists a neighborhood $V \ni x_0$, reduced representations $f_k = [f_k^0 : \dots : f_k^N]$ and **not necessarily reduced** representation $f = [f^0 : \dots : f^N]$ such that for every $0 \leq j \leq N$ the sequence f_k^j converge to f^j uniformly on V ;
- ii) $f_k \rightharpoonup f$ if and only if $f_k \xrightarrow{\Gamma} f$ and the limit representation $f = [f^0 : \dots : f^N]$ is **reduced**;
- iii) $f_k \rightarrow f$ if and only if $f_k \rightharpoonup f$ and corresponding **non-pluripolar** Monge-Ampère masses converge, i.e., for every $1 \leq p \leq n = \dim U$ one has

$$(dd^c \|z\|^2)^{n-p} \wedge (dd^c \ln \|f_k\|^2)^p \rightarrow (dd^c \|z\|^2)^{n-p} \wedge (dd^c \ln \|f\|^2)^p \quad (1.3)$$

weakly on compacts in U .

Here in (1.3) we suppose that $V = \Delta^n$, z, \dots, z_n are standard coordinates and $\|f\|^2 = |f^0|^2 + \dots + |f^N|^2$, i.e., $dd^c \ln \|f\|^2$ is the pullback of the Fubini-Study form by f . Non-pluripolar MA mass of $\ln \|f\|^2$ of order p here means

$$\int_{U \setminus Z_f} (dd^c \|z\|^2)^{n-p} \wedge (dd^c \ln \|f\|^2)^p, \quad (1.4)$$

where Z_f is the analytic sets of common zeroes of f^0, \dots, f^N . If this couple has no common divisors then $Z_f = I_f$, see section 4 for more details and the proof of this Theorem.

Remark 2. a) Reducibility or not of the limit representation $f = [f^0 : \dots : f^N]$ in this theorem doesn't depend on the choice of **converging** representations $f_k = [f_k^0 : \dots : f_k^N]$, provided they are taken to be reduced (the last can be assumed always). Indeed, any other reduced representation of f_k has the form $f_k = [g_k f_k^0 : \dots : g_k f_k^N]$, where g_k is holomorphic and nowhere zero. If the newly chosen representations converge to some representation of f then g_k must converge, say to g and this g is nowhere zero by Rouché's theorem. Therefore the obtained representation of the limit is $f = [g f^0 : \dots : g f^N]$ and it is reduced if and only if $f = [f^0 : \dots : f^N]$ was reduced.

b) The case when the representation $f = [f^0 : \dots : f^N]$ of the limit is not necessarily reduced was studied for mappings with values in \mathbb{P}^N by Fujimoto in [Fu], who called it **meromorphic**, or **m-convergence**. According to the part (i) of our theorem it turns out that our Γ -convergence (in the case of $X = \mathbb{P}^N$) is equivalent to **m-convergence** of Fujimoto.

1.2. Sets of normality and Bloch-Montel type criterion. In this paper we consider two classes of complex manifolds: projective and Gauduchon, the last is the class of complex manifolds carrying a dd^c -closed metric form - a Gauduchon form.

Let \mathcal{F} be a family of meromorphic mappings between complex manifolds U and X . \mathcal{F} is said to be *strongly/weakly* or *gamma* normal if from every sequence of elements of \mathcal{F} one can extract a subsequence converging on compacts in U in the corresponding sense. The maximal open subset $\mathcal{N}_{\mathcal{F}} \subset U$ on which \mathcal{F} is normal is called the set of normality. As it was already told the sets of strong normality could be arbitrary. In subsection 3.1 we prove the following

Theorem 3. *Let U be a domain in a Stein manifold \hat{U} such that \hat{U} is an envelope of holomorphy of U and let $f_k : \hat{U} \rightarrow X$ be a weakly converging on U sequence of meromorphic mappings with values in a disk-convex complex manifold X . Then:*

- (a) *If the weak limit f on f_k meromorphically extends from U to \hat{U} then f_k weakly converge to f on the whole of \hat{U} .*
- (b) *If, in addition, the manifold X carries a pluriclosed metric form then the weak limit f of f_k meromorphically extends to \hat{U} and then the part (a) applies.*

As a result the sets of weak normality are locally pseudoconvex provided the target is disk-convex and Gauduchon. Namely we have the following

Corollary 1. *Let $\mathcal{F} \subset \mathcal{M}(U, X)$ be family of meromorphic mappings from a complex manifold U to disk-convex Gauduchon manifold X . Then the set of weak normality $\mathcal{N}_{\mathcal{F}}$ of \mathcal{F} is pseudoconvex. If $\mathcal{F} = \{f_k\}$ is a sequence then the set of its convergence is pseudoconvex.*

Remark 3. The proof of this corollary clearly follows from Theorem 3. Sets of Γ -normality are also locally pseudoconvex under the same assumptions, see Proposition 3.1 in section 2.

As one more supporting argument in favor of weak convergence we prove in section 5 the following normality criterion

Theorem 4. *Let $\{H_i\}_{i=0}^d$, $d \geq 1$, be hypersurfaces in projective manifold X such that $Y := X \setminus \bigcup_{i=0}^d H_i$ is hyperbolically imbedded to X . Let \mathcal{F} be a family of meromorphic mappings from a complex manifold U to X such that:*

- i) *for every $i = 0, \dots, d$ and every compact $K \Subset U$ the volumes $f^*H_i \cap K$ counted with multiplicities are uniformly bounded for $f \in \mathcal{F}$;*
- ii) *\mathcal{F} uniformly separates every pair H_i, H_j , $0 \leq i < j \leq 2N$.*

Then the family \mathcal{F} is weakly normal on U .

Conditions (i) and (ii) are explained in section 5, they are intuitively clear and more or less necessary. The classically known case of a system of divisors with hyperbolically imbedded complement is $2N + 1$ hypersurfaces in \mathbb{P}^N in general position - Theorem of Bloch, see [Gr]. The case of m -normality (i.e., Γ -normality in our sense) was considered in [Fu].

1.3. Rational connectivity of the exceptional components of the limit. Strong convergence obviously implies the weak one:

$$\text{s-convergence} \implies \text{w-convergence}. \quad (1.5)$$

Our second principal task in this paper is to understand what obstructs a weakly converging sequence to converge strongly.

The problem is that by Theorem 1 the volumes of graphs of a strongly converging sequence are uniformly bounded over compacts in the source. When dimension n of the source U is two and X is Kähler the volumes of the graphs of a weakly converging sequence are bounded, see [Iv2]. The same is true if X is an arbitrary compact complex surface (and again $\dim U = 2$), see [Ne]. We shall say more about this in section 6.

But this turns out *not to be the case* starting from dimension three, i.e., the volumes of graphs of a weakly converging sequence can diverge to infinity over compacts of U . Via (1.3) this turns out to be a geometric counterpart of a well known discontinuity of Monge-Ampère masses, see Example 6.1 in section 4. Nevertheless for a sequence Γ_{f_k} of weakly converging meromorphic graphs we can consider the Hausdorff limit $\hat{\Gamma}$ (its always exists after taking a subsequence). Set $\Gamma := \hat{\Gamma} \setminus \Gamma_f$, where Γ_f is the graph of the limit map f , and call Γ a *bubble*. Set furthermore $\Gamma_a := \text{pr}_2(\text{pr}_1^{-1}(a) \cap \Gamma)$. We prove the following

Theorem 5. *Let X be a disk-convex Gauduchon manifold and let $f_k : U \rightarrow X$ be a weakly converging sequence of meromorphic mappings. Then for every point $a \in \Gamma$ the fiber Γ_a is either empty or is rationally connected.*

Here by saying that a closed subset Γ_a of a complex manifold is *rationally connected* we mean that every two distinct points $p, q \in \Gamma_a$ can be connected by a chain of rational curves, see section 7 for more details.

1.4. Dimension two and Fatou sets of meromorphic self-maps. Families of a special interest are the families of *iterates* $f^n := f \circ \dots \circ f$ of some fixed meromorphic self-map of a *compact* complex manifold X . The maximal open subset X where $\{f^n\}$ is relatively compact is called the Fatou set of f . Depending on the sense of convergence that one wishes to consider one gets different Fatou sets: strong, weak or gamma Fatou sets. We denote them as Φ_s , Φ_w and Φ_Γ respectively, their dependance on f will be clear from the context.

Corollary 2. *Let f be a meromorphic self map of a compact complex surface. Then the weak Fatou set Φ_w of f is locally pseudoconvex. If Φ_s is different from Φ_w then:*

- a) X is bimeromorphic to \mathbb{P}^2 ,
- b) $\Phi_w = X \setminus C$, where C is a rational curve in X ;
- c) the weak limit of any weakly converging subsequence $\{f^{n_k}\}$ of iterates is a degenerate map of X onto C .

It should be pointed out that our Fatou sets are different from the Fatou sets as they were considered in [FS]. In [FS] the Fatou set of f is the maximal open subset Φ of $X \setminus \overline{\bigcup_{n=0}^{\infty} f^{-n}(I_f)}$ where the family $\{f^n\}$ is equicontinuous (remark that on $X \setminus \overline{\bigcup_{n=0}^{\infty} f^{-n}(I_f)}$ all iterates are holomorphic). If, for example, $f : \mathbb{P}^2 \rightarrow \mathbb{P}^2$ is the Cremona transformation $[z_0 : z_1 : z_2] \rightarrow [z_1 z_2 : z_0 z_2 : z_0 z_1]$ then $\Phi_s = \Phi_w = \Phi_\Gamma = \mathbb{P}^2$ but $\Phi = \mathbb{P}^2 \setminus \{\text{three lines}\}$. In subsection 8.2 an example of higher degree and with interesting dynamics on the indeterminacy set is given. This is one more instance which shows how crucially can change a picture when the notion of convergence changes.

Note. Let us make a final note about the goals of this paper. We describe the most reasonable (on our opinion) notions of convergence of meromorphic mappings and conclude

that the *weak* one is the most appropriate. At the same time we detect that if a *weakly* converging sequence doesn't converge *strongly* then this imposes very serious restrictions on the target manifold (it is forced to contain many rational curves) and in some cases (ex. iterations) this puts strong constraints also on the sequence itself.

Acknowledgement. We are grateful to Alexander Rashkovskii for explaining to us the Example 6.1 with unbounded Monge-Ampère masses.

2. TOPOLOGIES ON THE SPACE OF MEROMORPHIC MAPPINGS

2.1. Complex manifolds and meromorphic mappings. Our manifolds will be Hausdorff and countable at infinity if the opposite is not explicitly stated. We shall also everywhere suppose that they are *disk-convex*.

Definition 2.1. A complex manifold X is called *disk-convex* if for every compact $K \Subset X$ there exists a compact \hat{K} such that for every $h \in \mathcal{O}(\Delta, X) \cap \mathcal{C}(\bar{\Delta}, X)$ such that $h(\partial\Delta) \subset K$ one has $h(\bar{\Delta}) \subset \hat{K}$.

\hat{K} is called the *disc envelope* of K . Let X be equipped with some Hermitian metric h . By ω_h denote the $(1,1)$ -form canonically associated with h . We say that the metric h is *d-closed* or *Kähler* if $d\omega_h = 0$. In the same way we say that h is *pluriclosed* or *Gauduchon* if $dd^c\omega_h = 0$. In [Ga] it was proved by P. Gauduchon that on a compact complex surface every Hermitian metric is conformally equivalent to the unique dd^c -closed one.

Remark 2.1. We shall need only existence of such metric forms on compact complex surfaces and this can be proved by duality: non existence of positive dd^c -closed $(1,1)$ -forms is equivalent to the presence of non-constant plurisubharmonic functions. The latter on a compact complex manifold is impossible.

We also fix some metric form ω_1 on U . In the case of a polydisc $U = \Delta^n$ we will work with the standard Euclidean metric e . The associated form will be denoted by $\omega_e = dd^c\|z\|^2 = \frac{i}{2} \sum_{j=1}^n dz_j \wedge d\bar{z}_j$. By $\text{pr}_1 : U \times X \rightarrow U$ and $\text{pr}_2 : U \times X \rightarrow X$ denote the projections onto the first and second factors. On the product $U \times X$ we consider the metric form $\omega = \text{pr}_1^*\omega_1 + \text{pr}_2^*\omega_h$.

A meromorphic mapping f between complex manifolds U and X is defined by an *irreducible* analytic subset $\Gamma_f \subset U \times X$ such that

- the restriction $\text{pr}_1|_{\Gamma_f} : \Gamma_f \rightarrow U$ of the natural projection to Γ_f is a proper modification, i.e., is proper and generically one to one.

Γ_f is called the *graph* of f . Due to the irreducibility of Γ_f and the Remmert proper mapping theorem the set of points over which pr_1 is not one to one is an analytic subset of U of codimension at least two. This set is called the set of points of indeterminacy of f and is usually denoted as I_f . Therefore another way to define a meromorphic mapping f between complex manifolds U and X is by considering a holomorphic map $f : U \setminus A \rightarrow X$, where A is an analytic subset of U of codimension at least two, such that the closure Γ_f of its graph is an analytic subset of the product $U \times X$ satisfying the condition above. Remark that the analyticity of the closure of the holomorphic graph is not automatic. Think about the natural projection $f : \mathbb{C}^2 \rightarrow \mathbb{C}^2 \setminus \{0\} / z \sim 2z$ of $\mathbb{C}^2 \setminus \{0\}$ onto a Hopf surface. The properness of the restriction of the projection pr_1 to the closure is, unless X is disc convex, not automatic to.

The volume of the graph Γ_f of a meromorphic mapping f is given by

$$\mathrm{Vol}(\Gamma_f) = \int_{\Gamma_f} \omega^n = \int_{\Gamma_f} (\mathrm{pr}_1^* \omega_1 + \mathrm{pr}_2^* \omega_h)^n = \int_U (\omega_1 + f^* \omega_h)^n, \quad (2.1)$$

where $n = \dim U$.

Remark 2.2. Let us make a few remarks concerning the notion of a meromorphic mapping.

a) If V is a subvariety of U such that $V \not\subset I_f$ then the restriction $f|_V$ of f to V is defined by taking as its graph $\Gamma_{f|_V}$ the *irreducible component* of the intersection $\Gamma_f \cap (V \times X)$ which projects onto V generically one to one. Therefore $\Gamma_{f|_V} \subset \Gamma_f \cap (V \times X)$ and the inclusion here is proper in general. The restriction of the graph to a (not necessarily a variety) $L \subset U$ is just the intersection $\Gamma_f \cap (L \times X)$.

In particular the *full image* of a set $L \subset U$ under f is defined as $f[L] := \mathrm{pr}_2(\Gamma_f \cap [L \times X])$.

b) It is probably worth to notice that $x \in I_f$ if and only if $\dim f[x] \geq 1$. This follows from the obvious observation that $I_f = \mathrm{pr}_1 \left(\{ (x_1, x_2) \in \Gamma_f : \dim_{(x_1, x_2)} \mathrm{pr}_1|_{\Gamma_f}^{-1}(x_1) \geq 1 \} \right)$.

c) If $\dim V = 1$ then the irreducible component of $\Gamma_f \cap (V \times X)$ which projects onto V is a curve. Since the projection is generically one to one it is one to one everywhere and therefore the restriction $f|_V$ is necessarily *holomorphic*.

d) Let us give the sense to $f^* \omega_h$ in the formula (2.1). The first integral there has perfectly sense since we are integrating a smooth form over a complex variety. Denote by I_f^ε the ε -neighborhood of the indeterminacy set I_f of f . Then (2.1) shows that the limit

$$\lim_{\varepsilon \rightarrow 0} \int_{U \setminus I_f^\varepsilon} (\omega_1 + f^* \omega_h)^n = \lim_{\varepsilon \rightarrow 0} \int_{U \setminus I_f^\varepsilon} \sum_{p=0}^n C_n^p \omega_1^{n-p} \wedge f^* \omega_h^p \quad (2.2)$$

exists. Therefore all $f^* \omega_h^p$ are well defined on U as positive currents.

2.2. Analytic cycles and currents. Before turning to the notions of convergence of meromorphic mappings let us recall the natural topologies on the space of analytic subsets of a complex manifold.

Recall that an analytic cycle of dimension r in a complex manifold Y is a formal sum $Z = \sum_j n_j Z_j$, where $\{Z_j\}$ is a locally finite sequence of reduced analytic subsets of pure dimension r and n_j are positive integers called multiplicities of Z_j . The set $|Z| := \bigcup_j Z_j$ is called the *support* of Z . All cycles in this paper, if the opposite is not stated, are supposed to have *connected* support. In our applications Y will be $U \times X$ and r will be the dimension $n = \dim U$.

By a coordinate chart adapted to Z we shall understand a relatively compact open set V in Y such that $V \cap |Z| \neq \emptyset$ together with a biholomorphism j of V onto a neighborhood \tilde{V} of $\bar{\Delta}^r \times \bar{\Delta}^q$ in \mathbb{C}^{r+q} , $r+q = \dim Y$, such that $j^{-1}(\bar{\Delta}^r \times \partial \Delta^q) \cap |Z| = \emptyset$. We shall denote such chart by (V, j) . The image $j(Z \cap V)$ of the cycle $Z \cap V$ under biholomorphism j is the image of the underlying analytic set together with multiplicities. Following Barlet and Fujiki, see [Ba] and [Fj], we call the quadruple $E = (V, j, \Delta^r, \Delta^q)$ a *scale* adapted to Z .

If $\mathrm{pr} : \mathbb{C}^r \times \mathbb{C}^q \rightarrow \mathbb{C}^r$ is the natural projection, then the restriction $\mathrm{pr}|_{j(Z \cap V)} : j(Z \cap V) \rightarrow \Delta^r$ is a branched covering of degree say d . This branched covering defines in a natural way a holomorphic mapping $\varphi_{j,Z} : \Delta^r \rightarrow \mathrm{Sym}^d(\Delta^q)$ - the d -th symmetric power of Δ^q

- by setting $\varphi_{j,Z}(z') = \{(\text{pr}|_{j(Z \cap V)})^{-1}(z')\}$ - unordered set of all preimages of z' under the projection in question. This construction, due to Barlet, allows to represent a cycle $Z \subset Y$ by a set of holomorphic maps $\varphi_{j_\alpha, Z} : \Delta^r \rightarrow \text{Sym}^d(\Delta^q)$, where $\{(V_\alpha, j_\alpha)\}$ is some open covering of $|Z|$ by adapted coordinate charts.

Definition 2.2. *One says that Z_k converges to Z in the topology of cycles if for every coordinate chart (V, j) adapted to Z there exists k_0 such that $\forall k \geq k_0$ this chart will be adapted to Z_k and the sequence of corresponding holomorphic mappings φ_{j, Z_k} converge to $\varphi_{j, Z}$ uniformly on Δ^r .*

This defines a metrizable topology on the space $\mathcal{C}_r(Y)$ of r -cycles in Y . This topology is equivalent to the **topology of currents**: $Z_k \rightarrow Z$ if for any continuous (r, r) -form χ with compact support one has

$$\int_{Z_k} \chi \rightarrow \int_Z \chi,$$

see [Fj]. It is also equivalent to the **Hausdorff** topology under an additional condition of **boundedness of volumes**. Recall that the Hausdorff distance between two subsets A and B of a metric space (Y, ρ) is a number $\rho(A, B) = \inf\{\varepsilon : A^\varepsilon \supset B, B^\varepsilon \supset A\}$. Here by A^ε we denote the ε -neighborhood of the set A , i.e. $A^\varepsilon = \{y \in Y : \rho(y, A) < \varepsilon\}$.

Now, $Z_k \rightarrow Z$ if and only if for every compact $K \Subset Y$ there exists $C_K > 0$ such that $\text{Vol}_{2r}(Z_k \cap K) \leq C_K$ and $Z_k \cap K \rightarrow Z \cap K$ with respect to the Hausdorff distance. This statement is the content of the Harvey-Shiffman's generalization of Bishop's compactness theorem. For the proof see [HS]. We denote the space of r -cycles on Y endowed with the topology described as above by $\mathcal{C}_r^{\text{loc}}(Y)$.

2.3. Strong convergence of meromorphic mappings. Let $\{f_k\}$ be a sequence of meromorphic mappings of a complex manifold U into a complex manifold X .

Definition 2.3. *We say that f_k converge **strongly** to a meromorphic map $f : U \rightarrow X$ (**s-converge**) if the sequence of graphs Γ_{f_k} converge over compacts to Γ_f in Hausdorff metric, i.e., for every compact $K \Subset U$ one has $\Gamma_{f_k} \cap (K \times X) \xrightarrow{H} \Gamma_f \cap (K \times X)$.*

Now let us prove Theorem 1 from Introduction, i.e., that Hausdorff convergence in the case of graphs implies the boundedness of volumes (over compacts) and therefore the convergence in the **topology of cycles**. Let us underline at this point that in this theorem one doesn't need to suppose anything on the target manifold X .

Proof of Theorem 1. The reason why Hausdorff convergence of graphs implies their stronger convergence in the topology of cycles is that, being the graphs, the analytic cycles Γ_{f_k} converge to Γ_f with **multiplicity** one. Now let us give the details. Let $a \in U \setminus I_f$ be a regular point of f and set $b = f(a)$. Then we can find neighborhoods $D_1 \ni a$ and $D_2 \ni b$ biholomorphic to Δ^n , $n = \dim U$ and Δ^p , $p = \dim X$ respectively such that $\Gamma_f \cap (\bar{D}_1 \times \partial D_2) = \emptyset$. In particular $V = D_1 \times D_2$ is an adapted chart for Γ_f and let $(V, j, \Delta^n, \Delta^p)$ be a corresponding scale. Here $j : V \rightarrow \Delta^n \times \Delta^p$ is some biholomorphism. From Hausdorff convergence of Γ_{f_k} to Γ_f we see that for $k \gg 1$ $\Gamma_{f_k} \cap (\bar{D}_1 \times \partial D_2) = \emptyset$. Therefore $\Gamma_{f_k} \cap (D_1 \times D_2) \rightarrow D_1$ is a ramified covering (i.e., is proper) of some degree d . But Γ_{f_k} is one to one over a generic point of D_1 . Therefore $d = 1$ and $\Gamma_{f_k} \cap V$ converge to $\Gamma_f \cap V$ as graphs (in particular as cycles). We proved that f_k converge to f on compacts of $U \setminus I_f$ as holomorphic mappings.

Let now $a \in I_f$ and take some $b \in f[a]$. As above take a neighborhood $V = D_1 \times D_2 \cong \Delta^n \times \Delta^p$ of (a, b) , where $a = 0$ and $b = 0$ in these coordinates. Denote by (w', w'') the coordinates in $\Delta^n \times \Delta^p$. Perturbing the slope of coordinate w'' we can suppose that $(\{0\} \times \Delta^p) \cap \Gamma_f$ has 0 as its isolated point.

Remark 2.3. After perturbation of the slope of w'' the decomposition $j(V) = \Delta^n \times \Delta^p$ will not correspond to the decomposition $U \times X$.

Now for sufficiently small $\varepsilon > 0$ we take as D_1 the disc Δ_ε^n , as D_2 the disc Δ_ε^p in perturbed coordinates \tilde{w}'' we get an adapted chart for Γ_f which possed the following property for the scale $E = (V, j, \Delta^n, \Delta^p)$:

- i) $\tilde{V} := j(V)$ writes as $\tilde{V} = \Delta^n \times \Delta^p$, $p = \dim X$.
- ii) Local coordinates (w', w'') of $\Delta^n \times \Delta^p$ enjoy the property that $z' := w' \in \Delta^n$ are local coordinates in U (but w'' is not a local coordinate on X).
- iii) $j(\Gamma_f \cap V) \rightarrow \Delta^n$ is a ramified covering of degree $d \geq 1$.

Again from Hausdorff convergence of Γ_{f_k} to Γ_f we get that for all $k \gg 1$ the intersection $j(\Gamma_{f_k} \cap V)$ is a ramified covering of Δ^n of degree d_k . Obviously $d_k \geq d$ for $k \gg 1$. If for some subsequence $d_k > d$ we shall get a contradiction as follows. In that case some irreducible component of $\Gamma_f \cap V$ will be approached by $\Gamma_{f_k} \cap V$ at least doubly. Let Γ stands for this irreducible component. Since $\dim [\Gamma_f \cap (I_f \times X)] \leq n - 1$ (by irreducibility of Γ_f) we see that Γ_{f_k} multiply approach every compact of $\Gamma \setminus (I_f \times X)$. Take a point $c \in \Gamma \setminus (I_f \times X)$ having a relatively compact neighborhood $W \subset \Gamma \setminus (I_f \times X)$ such that $\text{pr}_1|_W : W \rightarrow W_0$ is biholomorphic, i.e., W is the graph of f over $W_0 \Subset U \setminus I_f$. Now it is clear that $\Gamma_{f_k} \cap (W_0 \times X)$ cannot approach $\Gamma_f \cap (W_0 \times X) = W$ with multiplicity more than one, because Γ_{f_k} is a graph of a holomorphic map over W_0 for $k \gg 1$.

We proved that every irreducible branch of $\Gamma_f \cap V$ the graphs Γ_{f_k} approach with multiplicity one. Therefore $j(\Gamma_{f_k} \cap V) \rightarrow \Delta^n$ is a ramified covering of the same degree d as $j(\Gamma_f \cap V) \rightarrow \Delta^n$ for $k \gg 1$. This proves at a time that Γ_{f_k} converge to Γ_f in the topology of cycles and that their volumes are uniformly bounded. □

Strong convergence has some nice features, one was mentioned in the Introduction. Moreover, as it is explained in [Iv4], strong topology is natural in studying fix points of meromorphic self-mappings of compact complex manifolds. But domains of strong convergence and strong normality are quite arbitrary. We shall explain this in more details. Let \mathcal{F} be a family of meromorphic mappings from a complex manifold U to a disc convex complex manifold X .

Definition 2.4. *The set of normality of \mathcal{F} is the maximal open subset $\mathcal{N}_{\mathcal{F}}$ of U such that \mathcal{F} is relatively compact on $\mathcal{N}_{\mathcal{F}}$. If $\mathcal{F} = \{f_k\}$ is a sequence then the set of convergence of \mathcal{F} is the maximal open subset of U such that f_k converge on compacts of this subset.*

To be relatively compact in this definition means that from every sequence of elements of \mathcal{F} one can extract a converging on compacts subsequence. The sense of convergence (strong, weak or other) should be each time specified.

Example 2.1. 1. Let X be a Hopf three-fold $X := \mathbb{C}^3 \setminus \{0\}/z \sim 2z$. Denote by $\pi : \mathbb{C}^3 \setminus \{0\} \rightarrow X$ the canonical projection. Let $D \Subset \mathbb{C}^2$ be any bounded domain. Take a sequence $\{a_n\} \subset D$ accumulating to every point on ∂D . Let $g_n : \mathbb{C}^2 \rightarrow \mathbb{C}^3$ be defined as

$g_n(z) = (z, 1/n)$. Set $f_n := \pi \circ g_n$. Then the set of normality of $\mathcal{F} = \{f_n\}$ has D as one of its connected components.

Remark 2.4. For an analogous example with X projective see Example 4 from [Iv2].

2. The same example is instructive when understanding the notion of weak convergence. Take a converging to zero sequence a_n . Then f_n from this example will converge on compacts of $\mathbb{C}^2 \setminus \{0\}$ but the limit will not extend to zero meromorphically. I.e., f_n will not converge weakly in any neighborhood of the origin.

3. PSEUDOCONVEXITY OF SETS OF NORMALITY

3.1. Weak convergence and proof of Theorem 3. In view of such examples a weaker notion of convergence for meromorphic mappings was introduced in [Iv2]. Let $f \in \mathcal{M}(U, X)$ be a meromorphic map from U to X and let $\{f_k\} \subset \mathcal{M}(U, X)$ be a sequence of meromorphic mappings.

Definition 3.1. We say that f_k converge *weakly* to f (*w-converge*) if there exists an analytic subset A in U of codimension at least two such that f_k converge strongly to f on $U \setminus A$.

Remark 3.1. f_k converge weakly to f if and only if for every compact of $U \setminus I_f$ all f_k are holomorphic in a neighborhood of this compact for k big enough and uniformly converge there to f as holomorphic mappings. Indeed, let A be the minimal analytic set of codimension ≥ 2 such that f_k converge strongly to f on $U \setminus A$. Then A must be contained in I_f because if there exists a point $a \in A \setminus I_f$ then f is holomorphic in some neighborhood $V \ni a$ and then, by Rouché Principle of [Iv2] f_k for $k \gg 1$ are holomorphic on compacts in $V \setminus A$ and converge uniformly (on compacts) to f there. From here and the fact that $\text{codim } A \geq 2$ one easily gets that f_k are holomorphic on compacts in V and converge to f .

Now let us turn to the sets of weak convergence/normality. Sets of strong normality obviously are well defined, i.e., they do exist. The existence of sets of weak normality was proved in [Iv2], see Corollary 1.2.1a.

Remark 3.2. In the formulation of this Corollary the Author of [Iv2] speaks about “weak convergence” but the proof is about “weak normality”.

Domains of weak convergence of meromorphic mappings turn to be pseudoconvex for a large class of target manifolds. This follows from the “mutual propagation principle” stated in Theorem 3 in the Introduction. Let us give a proof of it.

Proof of Theorem 3. Let us prove the part (b) first.

Step 1. Extension of the limit. First of all by the main result of [Iv3] every meromorphic map $f : U \rightarrow X$ extends to a meromorphic map $f : \hat{U} \setminus A \rightarrow X$, where A is closed, complete $(n-2)$ -polar subset of \hat{U} . In more details that means that for every point $a \in A$ there exists a coordinate neighborhood $V \cong \Delta^{n-2} \times \mathbb{B}^2$ of $a = 0$ such that $A \cap (\Delta^{n-2} \times \partial \mathbb{B}^2) = \emptyset$ and for every $z' \in \Delta^{n-2}$ the intersection $A_{z'} := A \cap \mathbb{B}_{z'}^2$ is a zero dimensional complete pluripolar subset of $\mathbb{B}_{z'}^2 := \{z'\} \times \mathbb{B}^2$. Here \mathbb{B}^2 stands for the unit ball in \mathbb{C}^2 . Moreover, if $A \neq \emptyset$ then $f(\mathbb{S}_{z'}^3)$ is not homologous to zero in X . Here $\mathbb{S}_{z'}^3 = \partial \mathbb{B}_{z'}^2$ is the standard three-dimensional sphere in \mathbb{C}^2 .

Let U' be the maximal open subset of $\hat{U} \setminus (I_f \cup A)$ such that f_k converge to f on compacts in U' as holomorphic mappings.

Step 2. U' is locally pseudoconvex in $\hat{U} \setminus (I_f \cup A)$. If not there then by Docquier-Grauert criterion, see [DG], there would exist a point $b \in \partial U' \setminus (I_f \cup A)$ and a Hartogs figure $h : H_\varepsilon^n \rightarrow U'$ imbedded to U' such that the image $h(\Delta^n)$ of the corresponding polydisc contains b . All this is local and therefore we can assume that $h(\Delta^n)$ is relatively compact in $U \setminus (I_f \cup A)$. Pulling back f_k and f to Δ^n we arrive to contradiction as follows. Take a Stein neighborhood V of the graph of $f \circ h$ in $\Delta^n \times X$. Since for every compact $K \Subset H_\varepsilon^n$ we have that the graph of $f_k|_K$ is contained in V we conclude the same for every compact of Δ^n . Now $f_k \circ h$ converge to $f \circ h$ on compacts in Δ^n as holomorphic mappings. But that mean that they converge also around the preimage of b . Contradiction.

Since \hat{U} was supposed to be the envelope of holomorphy of U we obtain that $U' = \hat{U} \setminus (I_f \cup A)$.

Step 3. Removing A . Add I_f to A and take a sphere $\mathbb{S}_{z'}^3$ as described in **Step 1**. $f_k(\mathbb{S}_{z'}^3)$ is homologous to zero in X , because f_k meromorphically extends to the corresponding $\mathbb{B}_{z'}^2$. Moreover f_k converge to f in a neighborhood of $\mathbb{S}_{z'}^3$. This implies that $f_k(\mathbb{S}_{z'}^3)$ is also homologous to zero. Therefore A is empty, i.e., f is meromorphic on the whole of \hat{U} . The part (b) is proved.

The proof of (a) is a particular case of the **Step 2** of the proof of part (b). □

Remark 3.3. We gave a proof of Theorem 3 here because the proof of an analogous statement in [Iv2] uses a stronger extension claim from the subsequent paper [Iv3]. Namely the Author claimed that A appearing in the **Step 1** of the proof is *analytic* of codimension two. This was not achieved in [Iv3] (and is not clear for us up to know). Therefore we find necessary to remark that $(n-2)$ -polarity of A is, in fact, sufficient for this particular case.

3.2. Gamma convergence of meromorphic mappings. Let again f_k be a sequence of meromorphic mappings between complex manifolds U and X , the last is supposed to be disc convex. Let $f \in \mathcal{M}(U, X)$ be a meromorphic map.

Definition 3.2. We say that f_k Γ -converge to f if:

- i) there exists an analytic subset $A \subset U$ such that f_k strongly converge to f on compacts of $U \setminus A$;
- ii) for every divisor H in X , such that $f(U) \not\subset H$ and every compact $K \Subset U$ the volumes of $f_k^*H \cap K$ counted with multiplicities are uniformly bounded.

Remark 3.4. This notion is weaker than weak convergence because A can have components of codimension one. It might be convenient to add to A the indeterminacy set of f and then, see Remark 3.1, f_k will converge to f uniformly on compacts of $U \setminus A$ as holomorphic mappings. Condition (ii) is also satisfied for a weakly converging sequence, because divisors f^*H extend from $U \setminus A$ to U and if they have bounded volume on compacts of $U \setminus A$ then the same is true on compacts of U . All this obviously follows from the ingredients involved in the proof of Bishop's compactness theorem, see [St].

Example 3.1. a) Consider the following sequence of holomorphic mappings $f_k : \Delta \rightarrow \mathbb{P}^1$:

$$f_k : z \rightarrow \left[1 : 1 + \frac{1}{z} + \dots + \frac{1}{z^k k!} \right] = \left[z^k : z^k + z^{k-1} + \dots + \frac{1}{k!} \right]. \quad (3.1)$$

It is clear that f_k converges on compacts of $\Delta \setminus \{0\}$ to $f(z) = [1 : e^{\frac{1}{z}}]$ but, as it is clear from the second expression in (3.1) the preimage counting with multiplicities of the divisor $H = \{Z_0 = 0\}$ is $k[0]$ (here $[Z_0 : Z_1]$ are homogeneous coordinates in \mathbb{P}^1), i.e., has unbounded volume. And indeed, this sequence should not be considered as converging one, because its limit is not holomorphic on Δ .

b) Set $f_k(z) = [z : z - \frac{1}{k}] : \Delta \rightarrow \mathbb{P}^1$. This sequence clearly converges to the constant map $f(z) = [z : z] = [1 : 1]$ on compacts of $\Delta \setminus \{0\}$. Moreover, the preimage of any divisor $H = \{P(z_0, z_1) = 0\} \not\subset f_k(\Delta)$ is $\{z \in \Delta : P(z, z - \frac{1}{k}) = 0\}$, i.e., is a set of points, uniformly bounded in number counting with multiplicities. Therefore this sequence Γ -converge (but doesn't converge weakly).

Example 3.2. Consider the following sequence of meromorphic functions on Δ^2 (i.e., meromorphic mappings into \mathbb{P}^1):

$$f_k(z_1, z_2) = [z_1 : 2^{-k} z_2^k].$$

The limit map is constant $f(z) = [1 : 0]$. f_k converge to f strongly (uniformly in fact) on compacts of $\Delta^2 \setminus \{z_1 = 0\}$. If $[Z_0 : Z_1]$ are homogeneous coordinates in \mathbb{P}^1 then the preimage of the divisor $[Z_0 = 0]$ is $k[z_1 = 0]$, i.e., this sequence doesn't converge even in Γ sense.

Remark 3.5. Examples 3.1 and 3.2 are examples of converging outside of an analytic set of codimension one sequences which are not Γ -converging. In the first case the limit doesn't extend to the whole source, in the second it does. Convergence of meromorphic mappings of this type was introduced and studied by Rutishauser in [Ru].

If in Definition 2.4 the underlying convergence is Γ -convergence we get the corresponding notions of a convergence/normality set. Let us conclude this general discussion with the following

Proposition 3.1. *Let X be a disk-convex Gauduchon manifold. Then the sets of Γ -convergence/normality of meromorphic mappings with values in X are locally pseudoconvex.*

Proof. We shall prove the statement for the sets of Γ -normality, the case of sets of convergence obviously follows. Let D be the maximal open subset of U where the family \mathcal{F} is Γ -normal. Suppose that D is not pseudoconvex. Then by Docquier-Grauert criterion, see [DG], there exists an imbedding $h : H_\varepsilon^n \rightarrow D$ of a Hartogs figure into D such that h extends to an immersion of the polydisc to U with $h(\Delta^n) \cap \partial D \neq \emptyset$. Recall that Hartogs figure is the following domain

$$H_\varepsilon^n := (\Delta_\varepsilon^{n-1} \times \Delta) \cup (\Delta^{n-1} \times A_{1-\varepsilon,1}), \quad (3.2)$$

where $A_{1-\varepsilon,1} := \Delta \setminus \bar{\Delta}_{1-\varepsilon}$ is an annulus. Let us pull-back our family to Δ^n by h and therefore without loss of generality we can suppose that $U = \Delta^n$, \mathcal{F} is a family of meromorphic mappings from Δ^n to X , $H_\varepsilon^n \subset D \subset \Delta^n$ is the set of Γ -normality of \mathcal{F} such that $D \neq \Delta^n$.

That means that there exists a sequence $\{f_k\} \subset \mathcal{F}$, which converge on D but do not converge on compacts in Δ^n . Let us see that this is impossible. Let $f : D \rightarrow X$ be the

Γ -limit of f_k . Denote by A the analytic set in D such that f_k converge to f on compacts of $D \setminus A$. By [Iv3] f extends to $\Delta^n \setminus S$, where S is closed $(n-2)$ -complete polar subset of Δ^n . Let A' be the pure $(n-1)$ -dimensional part of A . By the theorem of Grauert we have two cases.

Case 1. The envelope of holomorphy of $D \setminus A'$ is Δ^n . In that case the Theorem 3 is applicable with $U = D \setminus A'$ and $\hat{U} = \Delta^n$ and gives us the weak (and therefore Γ) convergence of f_k on Δ^n .

Case 2. A' extends to a hypersurface \tilde{A} in Δ^n and $\Delta^n \setminus \tilde{A}$ is the envelope of holomorphy of $D \setminus A'$. In that case again by Theorem 3 f_k weakly converge to f on $\Delta^n \setminus \tilde{A}$. $S \setminus \tilde{A}$ is removable for f , see the Step 3 in the proof of Theorem 3. Therefore f_k strongly converge to f outside of a proper analytic set $A \cup I_f$. We need now to prove that f is extendable to Δ^n , i.e., that S is empty. By Proposition 2 of [Ne] the area of discs $f_k(\Delta_{z'})$ are bounded uniformly on k and on $z' \in \Delta^{n-1}(1-\varepsilon)$ for any fixed $\varepsilon > 0$, here $\Delta_{z'} := \{z'\} \times \Delta$. Therefore the areas of $f(\Delta_{z'})$ are bounded to. Theorem 1.5 together with Proposition 1.9 from [Iv3] imply now that f meromorphically extends onto $\Delta^{n-1}(1-\varepsilon) \times \Delta$. Therefore it extends to Δ^n . The condition (i) of Definition 3.2 is fulfilled.

Let H be a divisor in X . Then for every compact $K \Subset H_\varepsilon^n$ the volumes of $f_k^* H \cap K$ counted with multiplicities are bounded. By Oka-Riemanschnieder theorem, see [Rm], the volumes of the extensions of these divisors are bounded on compacts of Δ^n to. This verifies the condition (ii) of Definition 3.2. Proposition is proved. \square

4. CONVERGENCE OF MAPPINGS WITH VALUES IN PROJECTIVE SPACE

Now let us examine our notions of convergence on the example when the target manifold is a complex projective space.

4.1. Meromorphic mappings to complex projective space. Let a meromorphic mapping $f : U \rightarrow \mathbb{P}^N$ be given. Denote by $[w_0 : w_1 : \dots : w_N]$ the homogeneous coordinates of \mathbb{P}^N . Let $U_j = \{w \in \mathbb{P}^N : w_j \neq 0\}$ and let $\frac{w_0}{w_j}, \dots, \frac{w_N}{w_j}$ be affine coordinates in U_j . Set $D_j = f^{-1}(U_j)$. Since U_0 is isomorphic to \mathbb{C}^N the restriction $f|_{D_0} : D_0 \rightarrow U_0$ is given by holomorphic functions $\frac{w_1}{w_0} = f_1(z), \dots, \frac{w_N}{w_0} = f_N(z)$. The coordinate change in \mathbb{P}^N shows that $f|_{D_0 \cap D_j} : D_0 \cap D_j \rightarrow \mathbb{P}^N$ is given by functions $\frac{w_1}{w_0} = \frac{1}{f_j(z)}, \dots, \frac{w_N}{w_0} = \frac{f_N(z)}{f_j(z)}$ which are holomorphic in D_j . Therefore functions f_1, \dots, f_N are meromorphic on $D_0 \cup D_j$. This proves that f_1, \dots, f_N are meromorphic on $\cup_{j=0}^N D_j = U$. If $f_1 \equiv \dots \equiv f_N \equiv 0$ then $f(U) \equiv 0 \in U_0$. If not, let $f_1 \not\equiv 0$. One finds holomorphic functions h_j et g_j $0 \leq j \leq N$ in a polydisc neighborhood V of a given point $x \in U$, $g_j \neq 0$ such that

$$f_1 = \frac{h_1}{g_1}, \dots, f_N = \frac{h_N}{g_N}$$

and therefore gets

$$f := [1 : \frac{h_1}{g_1} : \dots : \frac{h_N}{g_N}] \prod_{j=1}^N g_j = [\prod_{j=1}^N g_j : h_1 \prod_{j=2}^N g_j : \dots : h_N \prod_{j=1}^{N-1} g_j].$$

This proves that f can be locally written in the form

$$f(z) := [f_0(z) : f_1(z) : \dots : f_N(z)] \quad (4.1)$$

as claimed.

4.2. Weak convergence of mappings with values in projective space. Let us prove now the part (ii) of Theorem 2 from the Introduction. I.e.,

Proposition 4.1. *A sequence of meromorphic mappings f_k from a complex manifold U to \mathbb{P}^N converges weakly on compacts of U if and only if for every point $z_0 \in U$ there exists a neighborhood $V \ni z_0$ and reduced representations $f_k = [f_k^0 : \dots : f_k^N]$, $f = [f^0 : \dots : f^N]$ in V such that for every $0 \leq j \leq N$ f_k^j converge to f^j uniformly on V .*

\Rightarrow Let $f_k \rightharpoonup f$, i.e., f_k converge to f weakly. Shrinking U we suppose that all f_k and f admit reduced representations

$$f_k = [f_k^0 : \dots : f_k^N] \quad (4.2)$$

and

$$f = [f^0 : \dots : f^N] \quad (4.3)$$

correspondingly. Up to making a linear coordinate change in \mathbb{P}^N we can suppose that $f[U]$ is not contained in any of coordinate hyperplanes, i.e., that $f^j \not\equiv 0$ for all $0 \leq j \leq N$. Set

$$Z^j = \{z \in U : f^j(z) = 0\},$$

and note that $\bigcap_{j=1}^N Z^j = I_f$. Since f_k converge on compacts in $U_j := U \setminus Z^j$ to f , see Remark 3.1, we see, taking $j = 0$, that

$$\frac{f_k^j}{f_k^0} \Rightarrow \frac{f^j}{f^0} \quad (4.4)$$

for all j on compacts in U_0 . Denote by Z_k^0 the zero divisors of f_k^0 and note that they leave every compact of U_0 as $n \rightarrow \infty$.

Lemma 4.1. *Divisors Z_k^0 converge to Z^0 in cycle space topology.*

Let us prove this Lemma first. Fix a point $a \in Z^0 \setminus Z^j$ (if $Z^0 \setminus Z^j$ is not empty) and take a relatively compact neighborhood $V \ni a$ such that $\bar{V} \cap Z^j = \emptyset$. We have that $f_k^0/f_k^j \Rightarrow f^0/f^j$ on \bar{V} . The Rouché's theorem easily implies now that $Z_k^0 \cap V$ converge to $Z^0 \cap V$ as currents.

Remark 4.1. In fact the cycle space topology on the space of divisors coincides with the topology of uniform convergence of defining them holomorphic functions, see [Stl]. And this immediately gives the previous assertion.

We conclude from here that Z_k^0 converge to Z^0 as cycles on compacts in $U \setminus I_f$. But then by [Ni], Theorem II, we obtain that they converge on the whole of U . Lemma 4.1 is proved.

We continue the proof of the Theorem. Shrinking U if necessary we can suppose that U is biholomorphic to $\Delta^n = \Delta^{n-1} \times \Delta$ and $Z_k^0 \cap U$ regularly covers Δ^{n-1} for $k \gg 1$. Now each Z_k^0 can be written as the zero set of a uniquely defined unitary polynomial P_k from $\mathcal{O}_{\Delta^{n-1}}[z_n]$ and these P_k uniformly converge to P - the defining polynomial for Z^0 . After multiplying each $[f_k^0 : \dots : f_k^N]$ by the unit P_k/f_k^0 we get the reduced representations

$$f_k = [P_k : g_k^1 : \dots : g_k^N].$$

The same with

$$f = [P : g^1 : \dots : g^N].$$

But now $P_k \rightrightarrows P$ and therefore from (4.4), which reads now as

$$\frac{g_k^j}{P_k} \rightrightarrows \frac{g^j}{P} \quad (4.5)$$

on compacts in U_0 , we get that for every $1 \leq j \leq N$ $g_k^j \rightrightarrows g^j$ on compacts in $U_0 = U \setminus Z^0$. But from the maximum principle it follows that $g_k^j \rightrightarrows g^j$ on compacts in U .

\Leftarrow For proving the inverse statement we start with converging reduced representations (4.2) to (4.3), i.e., $f_k^j \rightrightarrows f^j$ on U . Then for every $0 \leq j \leq N$ on every $U_j = U \setminus Z^j$ we get a convergence on compacts

$$\left(\frac{f_k^0}{f_k^j}, \dots, \frac{f_k^N}{f_k^j} \right) \rightrightarrows \left(\frac{f^0}{f^j}, \dots, \frac{f^N}{f^j} \right).$$

And since the codimension of $I_f = \bigcap Z^j$ is at least two we deduce the weak convergence of f_k to f . □

4.3. Strong convergence and convergence of meromorphic functions. Strong convergence of meromorphic maps into \mathbb{P}^N can be described in the following way. First, if $f_k \rightarrow f$ then $f_k \rightrightarrows f$. Therefore $[f_k^0 : \dots : f_k^N] \rightrightarrows [f^0 : \dots : f^N]$ for an appropriate reduced representations. According to (2.2) the volume of the graph of f_k is

$$\int_{U \setminus I_{f_k}} (\omega_1 + f_k^* \omega_{FS})^n = \int_{U \setminus I_f} \sum_{j=0}^n C_n^j \omega_1^j \wedge f_k^* \omega_{FS}^{n-j}. \quad (4.6)$$

Since $f_k^* \omega_{FS} = dd^c \ln \|f_k\|^2$ this is nothing but the non-pluripolar Monge-Ampère mass of $\ln \|f_k\|^2$ as appeared in (1.3). By Proposition 1 volumes of Γ_{f_k} converge to the volume of Γ_f , i.e.,

$$\int_{U \setminus I_{f_k}} \omega_1^j \wedge (dd^c \ln \|f_k\|^2)^{n-j} \rightarrow \int_{U \setminus I_f} \omega_1^j \wedge (dd^c \ln \|f\|^2)^{n-j} \quad (4.7)$$

for $0 \leq j < n$. In the case $U = \Delta^n$ this gives (1.3). Vice versa, if one has convergence of volumes the appearance of an exceptional component is impossible and we conclude the part (iii) of Theorem 2:

Proposition 4.2. *f_k converge to f strongly if and only if*

- i) *the appropriate reduced representations converge uniformly;*
- ii) *for every $0 \leq j \leq n-1$ one has (4.7).*

Now let us descend to the convergence of meromorphic functions. Meromorphic functions on a complex manifold U are exactly the meromorphic mappings from U to \mathbb{P}^1 . I.e., all our previous results and notions are applicable to this case.

Proposition 4.3. *If a sequence $\{f_k\}$ of meromorphic functions converge weakly then it converge strongly.*

Proof. Let f be the weak limit of f_k . We shall see in a moment, see Corollary 5.2 that volumes of graphs in this case are uniformly bounded over compacts in U . Therefore after going to a subsequence we get that the Hausdorff limit $\hat{\Gamma} := \lim \Gamma_{f_k}$ is a purely n -dimensional analytic subset of $U \times \mathbb{P}^1$. We claim that $\lim \Gamma_{f_k} = \Gamma_f$ in fact, i.e., that there are no exceptional components. If not take any irreducible component Γ of this limit different from Γ_f . Denote by γ its projection to U . γ is a proper analytic set of codimension at least two U . But then Γ should be contained in $\gamma \times \mathbb{P}^1$ and the last analytic set is of dimension $\dim U - 1$. This is impossible, because all components of $\lim \Gamma_{f_k}$ are of pure dimension $\dim U$. Therefore $\gamma = \emptyset$ and $\lim \Gamma_{f_k} = \Gamma_f$. □

4.4. Gamma convergence in projective case. In [Fu] and subsequent papers of Fujimoto the following type of convergence of meromorphic mappings with values in \mathbb{P}^N was considered, it was called the m -convergence (or *meromorphic* convergence): f_k m -converge to f if there exist reduced (admissible in the terminology of [Fu]) representations $f_k = [f_k^0 : \dots : f_k^N]$ which converge uniformly on compacts to $f = [f^0 : \dots : f^N]$, but the last *is not supposed* to be reduced (i.e., admissible), only not all f^j are identically zero. Let us prove the item (i) of Theorem 2.

Proposition 4.4. *When the target manifold X is the complex projective space \mathbb{P}^N the Γ -convergence of meromorphic mappings is equivalent to m -convergence in the sense of Fujimoto.*

Proof. \Rightarrow Suppose that $f_k \xrightarrow{\Gamma} f$. Let γ be the an analytic subset of U such that our sequence converge strongly on compacts of $U \setminus \gamma$. We add to γ also the indeterminacies of the limit f and therefore f_k will converge to f on $U \setminus \gamma$ in compact open topology. Let $f = [f^0 : \dots : f^N]$ be some reduced representation of the limit map. Making linear change of coordinates we can suppose, without loss of generality that $f^0 \not\equiv 0$, i.e., that $f(U) \not\subset H_0$, where $H_0 = \{Z_0 = 0\}$ in homogeneous coordinates $[Z_0 : \dots : Z_N]$ of \mathbb{P}^N . We have that $f_k^* H_0$ converge on compacts in U in the cycle space topology (after taking a subsequence).

Take some $a \in \gamma$ and choose a chart (V, j) adapted both to γ and $f^* H_0$ with coordinates z_1, \dots, z_n around a in such a way that $a = 0$ and $(\gamma \cup f^* H_0) \cap (\Delta^{n-1} \times \Delta)$ projects to Δ^{n-1} properly. Then $f_k^* H_0 \cap V$ also projects to Δ^{n-1} properly for $k \gg 1$. After going to a subsequence once more we can fix the degree d of ramified coverings $f_k^* H_0 \cap V \rightarrow \Delta^{n-1}$ and write the corresponding polynomials $P_k \in \mathcal{O}_{\Delta^{n-1}}[z_n]$ defining $f_k^* H_0 \cap V$. P_k converge to some P on (compacts of) Δ^{n-1} . Let $f_k = [f_k^0 : \dots : f_k^N]$ be some reduced representations of f_k on V . Notice that $f_k^* H_0 \cap V = \{f_k^0 = 0\}$. Divide each such representation by the unit f_k^0/P_k and get representations $f_k = [P_k : g_k^1 : \dots : g_k^N]$ with converging first terms P_k . At the same time $(g_k^1/P_k, \dots, g_k^N/P_k)$ represent f_k in nonhomogeneous coordinates of the chart $Z_0 \neq 0$ on \mathbb{P}^N . Therefore g_k^j/P_k converge to some f^j on compacts of $V \setminus (\gamma \cup f^* H_0)$. Therefore g_k^j converge to $g^j := f^j P$ on compacts of $V \setminus (\gamma \cup f^* H_0)$ to. By maximum principle they converge everywhere on V to the extension of g^j . We get that reduced representations $f_k = [P_k : g_k^1 : \dots : g_k^N]$ converge term by term to a (may be non reduced) representation $[P : g^1 : \dots : g^N]$ and this can be only a representation of f .

\Leftarrow Suppose now that f_k m -converge to f . Again change coordinates in \mathbb{P}^N , if necessary, in such a way that $f(U) \not\subset H_0$. Let V be a neighborhood of some point $a \in U$. If $a \in f^* H_0$ then take (V, j) to be an adapted chart to this divisor. In any case take V to be

biholomorphic to $\Delta^{n-1} \times \Delta$. Let $F_k = (f_k^0, \dots, f_k^N)$ be the lifts of f_k to \mathbb{C}^{N+1} in V such that F_k converge to the lift $F = (f^0, \dots, f^N)$ of f . From here one gets immediately that f_k^j/f_k^0 converge to f^j/f^0 uniformly on compacts of $V \setminus \{f^0 = 0\}$, i.e., that our maps converge strongly outside of a divisor.

Now let $H = \{P(Z_0, \dots, Z_N) = 0\}$ be a divisor such that $f(U) \not\subset H$. Using convergence of lifts $F_k = (f_k^0, \dots, f_k^N)$ to $F = (f^0, \dots, f^N)$ one gets that $f_k(U) \not\subset H$ for $k \gg 1$. One has also that $P(f_k^0, \dots, f_k^N)$ uniformly converge to $P(f^0, \dots, f^N)$ and this is equivalent to the convergence of divisors.

□

Remark 4.2. The relation between weak/gamma convergence and m -convergence for the case of $X = \mathbb{P}^N$ was indicated without proof in [Iv2].

5. BLOCH-MONTEL TYPE NORMALITY CRITERION

The aim of this section is to test the notion of weak convergence on the Bloch-Montel type normality statement, i.e., we are going to prove here the Theorem 4 from the Introduction.

5.1. Preliminaries. Before proceeding with the proof let us recall few basic facts. We start with an extended version of Zalcman's lemma, see [Me]:

Lemma 5.1. *A family \mathcal{F} of holomorphic mappings from Δ^n to a compact Hermitian manifold (X, h) is not normal at $z_0 \in \Delta^n$ if and only if there exist sequences $z_k \rightarrow z_0$, $r_k \searrow 0$, $f_k \in \mathcal{F}$ such that $f_k(z_k + r_k w)$ converge uniformly on compacts in \mathbb{C}^n to a nonconstant entire mapping $f : \mathbb{C}^n \rightarrow X$ such that $\|df(w)\|_h \leq 2$ for all $w \in \mathbb{C}^n$.*

This f may well have rank one. We shall also need the following result from [IS1], which is a precise version of Gromov compactness theorem (we shall need it in the integrable case only):

Proposition 5.1. *Let $u_k : \Delta \rightarrow X$ be a sequence of holomorphic maps into a disk-convex Hermitian manifold (X, h) with uniformly bounded areas, which uniformly converges on some annulus $A_{1-\varepsilon, 1}$ adjacent to the boundary $\partial\Delta$. Then u_k converge to stable complex curve over X after a reparametrization. Moreover, the compact components of the limit are rational curves.*

For the notions of stable curve over X , convergence after a reparametrization, as well as for the proof we refer to [IS1]. The obvious conclusion from this type of convergence is the following:

Corollary 5.1. *If u_k converge in stable sense to u and $u(\Delta)$ intersects a divisor H in X , but is not contained in H , then all $u_k(\Delta)$ intersect H for $k \gg 1$.*

Proof. It was proved in [IS2] (more details are given in [IS1]) that for any $k \gg 1$ one can join u_k with u by a holomorphic one parameter family of stable maps, see Proposition 2.1.3 in [IS2] for the exact statement. For us it is sufficient to understand that there exists a normal complex surface $Y \xrightarrow{\pi} \Delta$ foliated over the disc Δ such that all fibers $Y_s := \pi^{-1}(s)$ are discs and a holomorphic mapping $\mathcal{U} : Y \rightarrow X$ such that $\mathcal{U}|_{\Delta_0} = u$ and $\mathcal{U}|_{\Delta_{s_0}} = u_k$ for some $s_0 \in \Delta$.

Remark 5.1. The fact that this family can be contracted to a surface with normal points is proved in Lemma 2.2.6 in [IS2].

Let h be a defining holomorphic function of divisor H near the point of intersection $u(\Delta) \cap H$. Then $h \circ \mathcal{U}$ is holomorphic on Y (for this one might need to take disks of smaller radii) and is equal to zero at $0 \in Y_0 \subset Y$. At the same time it cannot vanish on $\bigcup_{s \in \Delta} \partial Y_s$ because $\mathcal{U}|_{Y_s}(\partial Y_s)$ is close to $u(\partial \Delta)$ for all $s \in \Delta$. Therefore the zero set of $h \circ \mathcal{U}$ must intersect every Y_s . And that means that $u_k(\Delta)$ intersects H . \square

Let us make one more remark. Let ω_{FS} be the Fubini-Study form on \mathbb{P}^N . For a holomorphic map $f : \bar{\Delta} \rightarrow \mathbb{C}^N$ (we always suppose f to be defined in a neighborhood of the closure $\bar{\Delta}$), the area of $f(\Delta)$ with respect to the Fubini-Study form is

$$\text{area}_{FS} f(\Delta) = \int_{\Delta} f^* \omega_{FS}. \quad (5.1)$$

Denote by $Z = (Z_0, \dots, Z_N)$ coordinates in \mathbb{C}^{N+1} and let $\pi : \mathbb{C}^{N+1} \rightarrow \mathbb{P}^N$ be the standard projection. Consider the following singular $(1,1)$ -form on \mathbb{C}^{N+1}

$$\omega_0 = dd^c \ln \|Z\|^2. \quad (5.2)$$

The following statement is a simple case of King's residue formula, but we shall give a simple proof for the sake of completeness.

Lemma 5.2. *For a holomorphic lift $F = (f^0, \dots, f^N) : \bar{\Delta} \rightarrow \mathbb{C}^{N+1}$ of $f : \bar{\Delta} \rightarrow \mathbb{P}^N$ (i.e., $f = \pi \circ F$) such that $F|_{\partial \Delta}$ doesn't vanishes one has*

$$\text{area}_{FS} f(\Delta) = \int_{\partial \Delta} d^c \ln \|F\|^2 - N_F. \quad (5.3)$$

Here N_F is the number of zeroes of F counted with multiplicities.

Proof. By the very definition of the Fubini-Study form one has $\pi^* \omega_{FS} = \omega_0$. And therefore it is immediate to check that in a neighborhood of a point $a \in \Delta$ such that $F(a) \neq 0$ one has that $f^* \omega_{FS} = F^* \omega_0$. As the result

$$\text{area}_{FS} f(\Delta) = \int_{\Delta} f^* \omega_{FS} = \int_{\Delta \setminus Z_F} F^* \omega_0, \quad (5.4)$$

where $Z_F := \{z_1, \dots, z_k\}$ is the set of zeroes of F , i.e., such z_l that $f^j(z_l) = 0$ for all $j = 0, \dots, N$. Let n_l be the multiplicity of zero z_l . Then $F(z) = (z - z_i)^{n_i} (g^0(z), \dots, g^N(z))$, where at least one of g^j -s is not zero at z_l . We have that

$$dd^c \ln \|F\|^2 = n_l \delta_{z_l} + dd^c \ln \|G\|^2,$$

where $G(z) = (g^0(z), \dots, g^N(z))$. Therefore $dd^c \ln \|G\|^2$ is an extension of $F^* \omega_0$ to z_l . The rest obviously follows from the Stokes formula. \square

Let us observe the following immediate corollary from this lemma.

Corollary 5.2. *Let $f_k : U \rightarrow \mathbb{P}^N$ be a Γ -converging sequence of meromorphic mappings and let L be a divisor in U such that f_k converge uniformly on compacts of $U \setminus L$. Let $V \cong \Delta^{n-1} \times \Delta$ be a chart adapted to L and to the limit M of $f_k^* H_0$, where $H_0 = [Z_0 = 0]$. Then the areas of the analytic discs $f_k(\Delta_{z'})$ are uniformly bounded in $z' \in \Delta^{n-1}$ and $k \in \mathbb{N}$.*

Proof. Let (z', z_n) be coordinates in $\Delta^{n-1} \times \Delta$. Denote by $F_k = (f_k^0, \dots, f_k^N)$ lifts of f_k to \mathbb{C}^{N+1} . Consider restrictions $f_k|_{\Delta_{z'}}$. Due to the fact that our chart is adapted to $M = \lim f_k^* H_0$ we have that f_k^0 doesn't vanish on $\partial\Delta_{z'}$ for $k \gg 1$ and, since it is also adapted to L the lifts $F_k = (f_k^0, \dots, f_k^N)$ converge in a neighborhood of $\partial\Delta_{z'}$. By (5.3) we have

$$\text{area}_{FS} f_k(\Delta_{z'}) \leq \int_{\partial\Delta_{z'}} d^c \ln \|F_k\|^2 \leq c, \quad (5.5)$$

i.e., the areas are uniformly bounded for $z' \in \Delta^{n-1}$ and all k .

□

Remark 5.2. For a family \mathcal{F} of meromorphic mappings from a manifold U to a projective manifold X to be normal an obvious necessary condition is that for any fixed hypersurface $H \subset X$ and any fixed compact $K \Subset U$ the volumes counting with multiplicities of intersections $f^*H \cap K$ should be uniformly bounded for $f \in \mathcal{F}$. It was proved by Fujimoto in [Fu] that this condition (in the case $X = \mathbb{P}^N$ and H_i are hyperplanes) turns out to be also sufficient, but only for the *meromorphic* normality. We in this paper are interested in the normality in the weak convergence sense (which is, that's to say, stronger than meromorphic). In that case there is one more necessary condition. Take two hypersurfaces H_0 and H_1 in X . Let $\{f^*H_0 : f \in \mathcal{F}\}$ and $\{f^*H_1 : f \in \mathcal{F}\}$ be the families of their preimages by elements of our family $f \in \mathcal{F}$. By boundedness of volumes condition for every sequence $f_{k_j}^* H_i$, $i = 0, 1$, some subsequences $f_{k_j}^* H_i$ converge to divisors L_0 and L_1 . If there exist *coinciding* (without taking account of multiplicities) components L'_0 and L'_1 of L_0 and L_1 respectively, then f_{k_j} cannot weakly converge in a neighborhood of $L'_0 = L'_1$. Indeed, since the limit f is a holomorphic map outside of I_f the preimages f^*H_0 and f^*H_1 cannot have common components. But L'_0 and L'_1 are such components. Contradiction. This will be formalized in the following definition.

Definition 5.1. We say that the family \mathcal{F} of meromorphic mappings from a complex manifold U to a complex manifold X uniformly separates hypersurfaces H_0 and H_1 from X if for any $f \in \mathcal{F}$ and any adapted for both f^*H_0 and f^*H_1 scale $E = (V, j, \Delta^{n-1}, \Delta)$ the Hausdorff distance between $f^*H_0 \cap V$ and $f^*H_1 \cap V$ for $f \in \mathcal{F}$ is bounded from below by a strictly positive constant.

Hausdorff distance is taken here in the Euclidean metric of \mathbb{C}^n . A constant in question may well depend on divisors H_0 , H_1 and adapted chart V , but it is supposed not to depend on $f \in \mathcal{F}$.

5.2. Proof of the normality criterion. Recall that a relatively compact open subset Y of a complex manifold X is said to be *hyperbolically imbedded* to X if for any two sequences $\{x_n\}$ and $\{y_n\}$ in Y converging to distinct points $x \in \bar{Y}$ and $y \in \bar{Y}$ one has

$$\limsup_{n \rightarrow \infty} k_Y(x_n, y_n) > 0,$$

where k_Y is the Kobayashi pseudodistance of Y . $Y \Subset X$ is said to be *locally hyperbolically complete* (l.h.c) if for every $y \in \bar{Y}$ there exists a neighborhood $V_y \ni y$ such that $V_y \cap Y$ is hyperbolically complete. For example every $Y \Subset X$ of the form $X \setminus \{\text{divisor}\}$ is obviously l.h.c. As it was proved in [Ki]: if Y is hyperbolically imbedded into X and is l.h.c. then Y is complete hyperbolic.

These notions are connected to complex lines in \bar{Y} by Theorem of Zaidenberg, see [Za]. By a complex line in Y (or in X) one understands an image of a non-constant holomorphic map $u : \mathbb{C} \rightarrow Y$ (or X). Sometimes one requires that $\|d_z u(\frac{\partial}{\partial x})\|_h \leq 1$ for all $z \in \mathbb{C}$, where h is some Hermitian metric on X . Complex line $u : \mathbb{C} \rightarrow \bar{Y} \Subset X$ is called *limiting for Y* if there exists a sequence of holomorphic mappings $u_n : \Delta(R) \rightarrow Y$ converging on compacts in \mathbb{C} to $u : \mathbb{C} \rightarrow \bar{Y}$. Theorem of Zaidenberg says now that: for a relatively compact l.c.h. domain Y in a complex manifold X to be complete hyperbolic and hyperbolically imbedded in X it is necessary and sufficient that Y doesn't contain complex lines and doesn't admits limiting complex lines.

Now we turn to the proof. Let $\{f_k\}$ be a sequence from \mathcal{F} , where \mathcal{F} satisfies the assumptions of Theorem 4 from the Introduction. $\{H_i\}_{i=0}^d$ our set of divisors.

Step 1. Convergence outside of a divisor. By Bishop's compactness theorem for every i some subsequence from $f_k^* H_i$ converges to a (may empty) hypersurface in U . Denote this limit hypersurface as L_i . Set

$$L := \bigcup_{i=0}^d L_i.$$

In order not to complicate notations we will not introduce subindexes when extracting subsequences.

If L is empty then for every compact $K \Subset U$ all f_k with k big enough send K to $X \setminus \bigcup_{i=0}^d H_i$, the last is Stein. In particular they are holomorphic in a neighborhood of K and we can use Zalcman's Lemma 5.1 together with Zaidenberg's characterization to extract a converging subsequence.

Therefore from now on we suppose L is nonempty. Take a point $z_0 \in U \setminus L$ and take a relatively compact neighborhood $V \ni z_0$ biholomorphic to a ball such that $\bar{V} \cap L = \emptyset$. Then for k big enough $f_k(\bar{V}) \subset X \setminus \bigcup_{i=0}^d H_i$. This implies that they all are holomorphic on V and we again can find a converging subsequence on V as before. Therefore some subsequence of $\{f_k\}$ (still denoted as $\{f_k\}$) converge on compacts of $U \setminus L$ in the usual sense of holomorphic mappings. Denote by f its limit. f is a holomorphic map from $U \setminus L$ to X .

Step 2. Convergence across the divisor. Take a point $z_0 \in L_0$, if L_0 is empty we can re-numerate L_i -s. fix an imbedding $i : X \rightarrow \mathbb{P}^N$ and let H be the intersection of X (i.e., of $i(X)$) with hyperplane $\{Z_0 = 0\}$ in the standard homogeneous coordinates $[Z_0 : \dots : Z_N]$ of \mathbb{P}^N . After going to a subsequence we have that $f_k^* H$ converge, denote by M the limit. Let (V, j) be an adapted chart for $L \cup M$ (and therefore also for L_0) at z_0 with the scale $E = (V, j, \Delta^{n-1}, \Delta)$. Let $P_k[z_n] \in \mathcal{O}_{\Delta^{n-1}}[z_n]$ be the defining unitary polynomial for $f_k^* H \cap V$. P_k converges to the defining polynomial P of $M \cap V$.

Let $f_k = [f_k^0 : \dots : f_k^N]$ be reduced representations of f_k on V (we write f_k for $i \circ f_k$). Then multiplying this representation by the unit P_k/f_k^0 we obtain a reduced representation $f_k = [P_k : g_k^1 : \dots : g_k^N]$. We have that g_k^j/P_k converge on compacts of $V \setminus (L \cup M)$. Therefore g_k^j converge there to, denote by g^j its limit. We see that lifts $F_k = (P_k, g_k^1, \dots, g_k^N)$ converge to $F := (P, g^1, \dots, g^N)$ on compacts of $V \setminus (L \cup M)$. By maximum principle they converge on V .

In particular f extends to a meromorphic mapping from U to X .

Remark 5.3. It is worth of noticing that at this stage we proved the Γ -normality of our family. For the case $X = \mathbb{P}^N$ with H_i hyperplanes this was proved in [Fu]. One more point worth of noticing is that the extendibility of f also follows from usual complex hyperbolic geometry, see [Ko].

Step 3. Convergence outside of codimension two. Changing indices of H_i , if necessary, we can suppose that our family uniformly separates H_0 and H_1 . Take a point $z_0 \in L_0 \setminus \bigcup_{i \neq 0} L_i$ such that L_0 in addition is smooth at z_0 . Take an adapted scale $E = (V, j, \Delta^{n-1}, \Delta)$ for L_0 near z_0 which intersects L only by the smooth part of L_0 and, moreover, such that $j(L_0 \cap V) = d \cdot [\Delta^{n-1} \times \{0\}]$ for some multiplicity $d \geq 1$. Fix coordinates (z', z_n) on $\Delta^{n-1} \times \Delta$. By Corollary 5.2 the areas of analytic discs $f_k|_{\Delta_{z'}}$ are uniformly bounded. Fix some $z' \in \Delta^{n-1}$ and take a subsequence f_k such that $f_k|_{\Delta_{z'}}$ converge in stable topology to $f|_{\Delta_{z'}}$ plus a chain $C_{z'}$ of rational curves. By Corollary 5.1 if $C_{z'}$ intersects some H_1 with $i \neq 0$ then $f_k|_{\Delta_{z'}}(\Delta_{z'})$ intersects H_1 to. But then $f_k^* H_1 \cap V$ is nonempty and converge to $L_1 \cap V$. This can be only $L_0 \cap V$ with some multiplicity, because V was chosen in such a way that $L \cap V = L_0 \cap V$. The last violates the assumed uniform separability of the pair H_0, H_1 by \mathcal{F} . Therefore $C_{z'}$ is empty. That means that (some subsequence of) $f_k|_{\Delta_{z'}}$ uniformly on $\Delta_{z'}$ converge to $f|_{\Delta_{z'}}$. This implies that the whole sequence f_k restricted to $\Delta_{z'}$ converge to f . Therefore f_k converge to f on $U \setminus \text{Sing } L$ in compact open topology as holomorphic mappings. This proves the Theorem. \square

Remark 5.4. Theorem of Bloch, see also [Gr], states that $Y = \mathbb{P}^N \setminus \bigcup_{j=0}^{2N} H_j$ is hyperbolically imbedded to \mathbb{P}^N , where H_i are hyperplanes in general position. Therefore $Y = \mathbb{P}^N \setminus \bigcup_{i=0}^{2N} H_i$ is an example for our Theorem 4.

6. BEHAVIOR OF VOLUMES OF GRAPHS UNDER WEAK AND GAMMA CONVERGENCE

In this section we are concerned with the following question: let meromorphic mappings $f_k : U \rightarrow X$ converge in some sense to a meromorphic map f , what can be said about the behavior of volumes of graphs of f_k over compacts in U ? If f_k converge to f strongly then, as it was proved in Theorem 1, for every relative compact $V \Subset U$ we have that

$$\text{Vol}(\Gamma_{f_k|_V}) \rightarrow \text{Vol}(\Gamma_{f|_V}). \quad (6.1)$$

When f_k converges only weakly one cannot, of course expect anything like (6.1). At most what one can expect is that volumes of Γ_{f_k} stay bounded over compacts in U and converge to the volume of Γ_f plus volumes of exceptional components. I.e., the question is if for a weakly converging sequence $\{f_k\}$ one has that for every relatively compact open $V \Subset U$ there exists a constant C_V such that

$$\text{Vol}(\Gamma_{f_k|_V}) \leq C_V \quad \text{for all } k. \quad (6.2)$$

This turns to be wrong in general, the following example was communicated to us by A. Rashkovskii.

6.1. Example of Rashkovskii.

Example 6.1. There exists a sequence $\varepsilon_k \searrow 0$ such that *holomorphic* mappings $f_k : \mathbb{B}^3 \rightarrow \mathbb{P}^3$ defined as

$$f_k : (z_1, z_2, z_3) \rightarrow [z_1 : z_1 - \varepsilon_k : z_2 : z_3^k] \quad (6.3)$$

converge weakly to $f(z) = [z_1 : z_1 : z_2 : 0]$ on compacts of the unit ball $\mathbb{B}^3 \subset \mathbb{C}^3$, but the volumes of graphs of f_k over the ball $\mathbb{B}^3(1/2)$ of radius $1/2$ diverge. In fact

$$\text{Vol}(\Gamma_{f_k}) \cap (\mathbb{B}^3(1/2) \times \mathbb{P}^3) \geq k. \quad (6.4)$$

Consider the following family of plurisubharmonic functions on the unit ball \mathbb{B}^3 in \mathbb{C}^3 :

$$u_{\varepsilon,k}(z) = \ln(|z_1|^2 + |z_1 - \varepsilon|^2 + |z_2|^2 + |z_3|^k), \quad \varepsilon \in (0, 1/4). \quad (6.5)$$

Note that every $u_{\varepsilon,k}$ is bounded in \mathbb{B}^3 and its total MA mass in $\mathbb{B}^3(1/2)$ coincides with those of the function

$$\tilde{u}_{\varepsilon,k} := \max\{u_{\varepsilon,k}, s_k\} \quad \text{where} \quad s_k = \min\{\ln|z_1|^2 + |z_2|^2 + |z_3|^k : z \in \mathbb{S}^5(1/2)\}.$$

Here $\mathbb{S}^5(1/2) = \partial\mathbb{B}^3(1/2)$ is the sphere of radius $1/2$. This fact follows from the Bedford-Taylor definition of the MA mass of a product of *bounded* psh functions, see [BT]: $dd^c u_1 \wedge dd^c u_2 := dd^c(u_1 dd^c u_2)$ and so on by induction. Here the point is, of course, to prove that $dd^c(u_1 dd^c u_2)$ is again a closed positive current. Now one writes

$$\begin{aligned} \text{MA}_{\mathbb{B}^3(1/2)}(u_{\varepsilon,k}) &= \int_{\mathbb{B}^3(1/2)} (dd^c u_{\varepsilon,k})^3 = \int_{\mathbb{B}^3(1/2)} dd^c u_{\varepsilon,k} \wedge (dd^c u_{\varepsilon,k})^2 = \int_{\partial\mathbb{B}^3(1/2)} d^c u_{\varepsilon,k} \wedge (dd^c u_{\varepsilon,k})^2 = \\ &= \int_{\partial\mathbb{B}^3(1/2)} d^c \tilde{u}_{\varepsilon,k} \wedge (dd^c \tilde{u}_{\varepsilon,k})^2 = \int_{\mathbb{B}^3(1/2)} (dd^c \tilde{u}_{\varepsilon,k})^3 = \text{MA}_{\mathbb{B}^3(1/2)}(\tilde{u}_{\varepsilon,k}) \end{aligned}$$

because $u_{\varepsilon,k} = \tilde{u}_{\varepsilon,k}$ on the sphere $\mathbb{S}^5(1/2)$. Since $\tilde{u}_{\varepsilon,k}$ converge uniformly to $\tilde{u}_k = \max\{\ln(2|z_1|^2 + |z_2|^2 + |z_3|^k), s_k\}$ as $\varepsilon \searrow 0$ and $\text{MA}_{\mathbb{B}^3(1/2)}(\tilde{u}_k) = \text{MA}_{\mathbb{B}^3(1/2)}(u_k) = 4k$, where $u_k = \ln(2|z_1|^2 + |z_2|^2 + |z_3|^k)$, we shall have that for ε_k small enough $\text{MA}_{\mathbb{B}^3(1/2)}(u_{\varepsilon_k,k}) \geq k$. This finishes the proof.

Remark 6.1. Examples of psh functions with polar singularities and unbounded non-polar MA mass were constructed first by Shiffman and Taylor, see [Si], and especially simple one by Kiselman, see [Ks]: $u(z_1, \dots, z_n) = (1 - |z_n|^2)(-\ln \|z'\|^2)^{1/2}$ for $z' = (z_1, \dots, z_{n-1})$. Taking any of these examples and smoothing it by convolutions one gets a decreasing sequences of psh functions converging outside of an analytic set (on any codimension) to a psh function, smooth outside of this set with unbounded non-polar MA mass. The remarkable feature of the example of Rashkovskii, just described, is that functions in this example have a *geometric meaning*, their dd^c -s are pullbacks of Fubini-Study from by a meromorphic mappings to the complex projective space, i.e., the sum of their non-polar MA masses are the volumes of the corresponding graphs.

6.2. Case of dimensions one and two. If $\{f_k\}$ is a gamma-converging sequence of meromorphic mappings with values in one dimensional complex manifold then it is easy to see that areas of graphs of f_k -s are locally bounded over compacts in the source. Indeed, a one dimensional manifold X either properly imbeds to \mathbb{C}^n (when X is noncompact) or is projective and therefore imbeds to \mathbb{P}^n . In both cases by Theorem 2 we have convergence of reduced representations to a, may be nonreduced representation of the limit. Inequality (5.5) implies that in an appropriately chosen local coordinates (z', z_n) one has

$$\text{Vol}(\Gamma_{f_k|_{\Delta^n}}) = \int_{\Delta^n} (dd^c |z|^2)^n + \int_{\Delta^n} (dd^c |z|^2)^{n-1} \wedge f_k^* \omega_{FS} \leq \int_{\Delta^n} (dd^c |z|^2)^n +$$

$$+ \int_{\Delta^{n-1}} (dd^c |z|^2)^{n-1} \int_{\partial \Delta_{z'}} d^c \ln \|F_k\|^2 \leq \text{const.}$$

Next, if the dimension n of the source U is 2 the boundedness of volumes of graphs of a weakly converging sequence is automatic. This can be seen at least in two ways.

First, in projective case this readily follows from the following formula of King, see [Kg]:

$$d [d^c \ln(\|f\|^2) \wedge dd^c \ln(\|f\|^2)] = \chi_{U \setminus I_f} \left[(dd^c \ln(\|f\|^2))^2 \right] - \sum_j n_j [Z_j], \quad (6.6)$$

provided I_f has pure codimension two. Z_j are irreducible components (branches) of the indeterminacy set $I_f = Z_f$ of f . If it has branches of higher codimension then around these branches a higher order non-pluripolar masses can be expressed in a similar way. Now if f_k weakly converge to f formula (6.6) immediately gives a uniform bound of corresponding MA masses (even together with that concentrated on pluripolar sets I_{f_k}). If $n = 2$ then that's all we need.

Second, using Skoda potentials, or Green functions, as it was done in [Iv2] Theorem 2, one can bound non-pluripolar Monge-Ampère masses of order two also in the case of weakly converging sequence with values in disk-convex Kähler X . This observation implies that if X is disk-convex Kähler and $\dim U = 2$ then the volumes of graphs of weakly converging sequences of meromorphic mappings $U \rightarrow X$ are uniformly bounded over compacts in U .

Moreover, it was proved in [Ne] that volumes of weakly converging sequence are bounded also in the case when X is any compact complex surface. The proof uses Kähler case separately and then the fact that a non-Kähler surface has only finitely many rational curves.

Remark 6.2. Let us remark that there is one more important case when the volumes of graphs of weakly (even Γ) converging sequence necessarily stay bounded: namely when $\{f_k\}$ is a Γ -converging sequence of meromorphic mappings between projective manifolds X and Y . Indeed the volumes of graphs Γ_{f_k} are uniformly bounded as it is straightforward from Besout theorem.

7. RATIONAL CONNECTIVITY OF THE EXCEPTIONAL COMPONENTS OF THE LIMIT

7.1. Chains of rational curves and proof of Theorem 5. Recall that a rational curve C in a complex manifold (or space) X is an image of \mathbb{P}^1 in X under a nonconstant holomorphic map $h : \mathbb{P}^1 \rightarrow X$. A chain of rational curves is a *connected* union $C = \bigcup_j C_j$ of finitely many rational curves.

Definition 7.1. A closed subset $\Gamma \subset X$ we call *rationally connected* if for very two points $p \neq q$ in Γ there exists a chain of rational curves $C \subset \Gamma$ such that $p, q \in C$.

One says also that C *connects* p with q . If Γ is a complex manifold then this property is equivalent to the either of the following two ones:

- Every two points in X can be connected by a single rational curve.
- For any finite set of points $F \subset X$ there exists a rational curve $C \supset F$.

We refer to [Ar] for these facts. Now let us turn to the proof of Theorem 5 from the Introduction. It consists from the two following lemmas.

Lemma 7.1. *Let $V \cong \Delta^{n-1} \times \Delta$ be a chart adapted to γ . Suppose that there exists a dense subset $S \subset \Delta^{n-1}$ such that the areas of the analytic discs $\Gamma_{f_k|_{\Delta_{z'}}$ are uniformly bounded in $z' \in S$ and $k \in \mathbb{N}$ then for every $a \in \gamma$ the fiber $\Gamma_a := \text{pr}_2(\text{pr}_1^{-1}(a))$ is rationally connected.*

Proof. here writing $f_k|_{\Delta_{z'}}$ we mean the restriction of f_k to the disk $\Delta_{z'} := \{z'\} \times \Delta$. Fix a point $a \in \gamma$ and some $a^1, a^2 \in \Gamma_a$. Suppose $a^1 \neq a^2$, otherwise there is nothing to prove. We need to prove that there exists a chain of rational curves in Γ_a connecting a^1 with a^2 . Since Γ_{f_k} converge to $\hat{\Gamma} \supset \Gamma_a$ there exist $a_k^1 \rightarrow a$ and $a_k^2 \rightarrow a$ such that $f_k(a_k^1) \rightarrow a^1$ and $f_k(a_k^2) \rightarrow a^2$. Perturbing slightly we can take such a_k^i to be regular (i.e., not indeterminacy) points of f_k for $i = 1, 2$. Take a scale adapted to γ near a in the sense that $\gamma \cap (\Delta^{n-1} \times \partial\Delta) = \emptyset$. Denote by $(z', z_n) = (z_1, \dots, z_{n-1}, z_n)$ the corresponding coordinates and assume without loss of generality that $a = 0$.

Let $b_k^1 \rightarrow 0'$ and $b_k^2 \rightarrow 0'$ in Δ^{n-1} be such that $a_k^1 \in \Delta_{b_k^1}$ and $a_k^2 \in \Delta_{b_k^2}$. Taking again a_k^i sufficiently general we can arrange that $b_k^i \in S$ and discs $\Delta_{b_k^1}$ and $\Delta_{b_k^2}$ are converge to the disc $\Delta_{0'}$.

After taking a subsequence we get that graphs in question converge to the graph $\Gamma_{f|_{\Delta_{0'}}} \cup C^i$, where $C^i \subset \{a\} \times X$ are chains of rational curves. Both these chains contain the point $f|_{\Delta_{0'}}(a)$. Therefore $C := C^1 \cup C^2$ is connected. At the same time by construction $C^i \ni a^i$. The lemma is proved. □

The second lemma is the following

Lemma 7.2. *Let \mathcal{F} be a family of meromorphic mappings from Δ^n to a disk-convex manifold X , which admits a pluriclosed metric form. Suppose that for some $0 < \epsilon < 1$, the family \mathcal{F} is holomorphic and equicontinuous on the Hartogs figure H_ϵ^n . Then for every $0 < r < 1$ areas of graphs $\Gamma_{f_{z'}}$ of restrictions $f_{z'} := f|_{\Delta_{z'}(r)}$ of $f \in \mathcal{F}$ to the discs $\Delta_{z'}(r) := \{z'\} \times \Delta_r$ are uniformly bounded in $z' \in \Delta_r^{n-1}$ and $f \in \mathcal{F}$.*

Proof. For $f : \Delta^n \rightarrow X$ a meromorphic map, we denote by $I_f \subset \Delta^n$ the set of points of indeterminacy of f . Since we suppose that all $f \in \mathcal{F}$ are holomorphic on H_ϵ^n the sets I_f do not intersect $\Delta^{n-1} \times A_{1-\epsilon, 1}$. Consider currents $T_f = f^*\omega$ on Δ^n , where ω is a pluriclosed metric form on X . Write

$$T_f = \frac{i}{2} t_f^{\alpha\bar{\beta}} dz_\alpha \wedge d\bar{z}_\beta,$$

where $t_f^{\alpha\bar{\beta}}$ are distributions on Δ^n (in fact measures), smooth on $\Delta^n \setminus I_f \supset H_\epsilon^n$. Fix $1 - \epsilon < r < r_1 < 1$ and consider on $\Delta^{n-1} \setminus \pi(I_f)$ (where $\pi : \Delta^n \rightarrow \Delta^{n-1}$ is the canonical projection onto the first factor) the **area functions** a_f given by

$$a_f(z') = \text{area } f_{z'}(\Delta_{r_1}) = \int_{\Delta_{z'}(r_1)} T_f = \frac{i}{2} \int_{\Delta_{z'}(r_1)} t_f^{n\bar{n}} dz_n \wedge d\bar{z}_n. \quad (7.1)$$

Functions a_f are well-defined and smooth on $\Delta^{n-1} \setminus \pi(I_f)$.

The proof of Proposition will be done in two steps.

Step 1. *Distributions $t_f^{n\bar{n}}$ are locally integrable in Δ^n .* Note that forms T_f are smooth on H_ϵ^n and the family $\{T_f : f \in \mathcal{F}\}$ is equicontinuous there. The condition that $dd^c T_f = 0$

implies, in particular, that for all $1 \leq k, l \leq n-1$ one has

$$\frac{\partial^2 t_f^{n\bar{n}}}{\partial z_k \partial \bar{z}_l} + \frac{\partial^2 t_f^{k\bar{l}}}{\partial z_n \partial \bar{z}_n} - \frac{\partial^2 t_f^{k\bar{n}}}{\partial z_n \partial \bar{z}_l} - \frac{\partial^2 t_f^{n\bar{l}}}{\partial z_k \partial \bar{z}_n} = 0. \quad (7.2)$$

From (7.2) we get that on $\Delta^{n-1} \setminus \pi(I_f)$:

$$\begin{aligned} dd^c a_f &= \left(\frac{i}{2}\right)^2 \sum_{k,l=1}^{n-1} \left(\int_{\Delta_{z'}(r_1)} \frac{\partial^2 t_f^{n\bar{n}}}{\partial z_k \partial \bar{z}_l} dz_n \wedge d\bar{z}_n \right) dz_k \wedge d\bar{z}_l = \\ &= \left(\frac{i}{2}\right)^2 \sum_{k,l=1}^{n-1} \left(\int_{\Delta_{z'}(r_1)} \frac{\partial^2 t_f^{k\bar{n}}}{\partial z_n \partial \bar{z}_l} + \frac{\partial^2 t_f^{n\bar{l}}}{\partial z_k \partial \bar{z}_n} - \frac{\partial^2 t_f^{k\bar{l}}}{\partial z_n \partial \bar{z}_n} \right) dz_n \wedge d\bar{z}_n \cdot dz_k \wedge d\bar{z}_l = \\ &= \left(\frac{i}{2}\right)^2 \sum_{k,l=1}^{n-1} \left(\int_{\partial \Delta_{z'}(r_1)} \frac{\partial t_f^{k\bar{n}}}{\partial \bar{z}_k} d\bar{z}_n + \int_{\partial \Delta_{z'}(r_1)} \frac{\partial t_f^{n\bar{l}}}{\partial z_k} dz_n - \int_{\partial \Delta_{z'}(r_1)} \frac{\partial t_f^{k\bar{l}}}{\partial \bar{z}_n} d\bar{z}_n \right) dz_k \wedge d\bar{z}_l =: \phi_f. \end{aligned} \quad (7.3)$$

Forms ϕ_f are smooth in the whole unit polydisc Δ^{n-1} and equicontinuous there because forms T_f are smooth in $\Delta^{n-1} \times A_{1-\varepsilon,1} \subset H_\varepsilon^n$ and equicontinuous there. Let us find a smooth and equicontinuous family on Δ_r^n of solutions ψ_f of

$$dd^c \psi_f = \phi_f. \quad (7.4)$$

Set

$$h_f := a_f - \psi_f. \quad (7.5)$$

Since a_f is positive on $\Delta^{n-1} \setminus \pi(I_f)$ and ψ_f is smooth on Δ^{n-1} we see that h_f is bounded on Δ^{n-1} from below. Also $dd^c h_f = 0$ on $\Delta^{n-1} \setminus \pi(I_f)$ and therefore h_f extends to a plurisuperharmonic function on Δ^{n-1} . This implies that $h_f \in L_{loc}^1(\Delta^{n-1})$ see [Ho]. It follows that a_f and $t_f^{n\bar{n}}$ are locally integrable. Step 1 is proved.

Step 2. Under the hypotheses of Lemma 7.2 functions a_f defined by (7.1) are smooth on Δ^{n-1} and for every fixed $r < 1$ the family $\{a_f\}_{f \in \mathcal{F}}$ is equicontinuous on $\bar{\Delta}_r^{n-1}$. Function h_f given by (7.5) is plurisuperharmonic in Δ^{n-1} and pluriharmonic on $\Delta^{n-1} \setminus \pi(I_f)$. Therefore by Siu's lower semicontinuity of the level sets of Lelong numbers we have

$$dd^c h_f = - \sum_{A \text{ irr.comp. of } \pi(I_f)} c_A(f)[A], \quad (7.6)$$

where $c_A(f) \geq 0$ and $[A]$ denotes the current of integration over the irreducible component A of $\pi(I_f)$ of codimension one.

Remark 7.1. Note that through components of higher codimension a pluriharmonic function h_f extends (as a pluriharmonic function). Therefore in (7.6) the sum is taken over the components of codimension one only.

We need to prove that $c_A(f) = 0$. From (7.5) we get

$$dd^c a_f = dd^c \psi_f - \sum_{A \text{ irr.comp. of } \pi(I_f)} c_A(f)[A], \quad (7.7)$$

where dd^c from a_f is taken in the sense of distributions (as from L^1_{loc} -function). Let $\{h_A\}$ be equations of A . By Poincaré formula, see [GK], $[A] = dd^c \ln |h_A|^2$ and therefore (7.7) writes as

$$dd^c a_f = dd^c \psi_f - \sum_{A \text{ irr.comp. of } \pi(I_f)} c_A(f) dd^c \ln |h_A|^2 \quad (7.8)$$

Take an one dimensional disk Δ in Δ^{n-1} which intersects $\pi(I_f)$ transversely at points $\{z_j\}$. Then (7.7) gives for restrictions of a_f and φ_f to Δ (and we shall denote them by the same letters) the following

$$\Delta a_f = \Delta \psi_f - \sum_{z_j \in \pi(A_f)} c_j(f) \delta_{z_j(f)}. \quad (7.9)$$

Fix $\delta > 0$ such that $\Delta(\delta, z_j)$ are pairwise disjoint. Let φ be a test function on Δ with support in $\Delta(\delta, z_j)$ for some fixed j . The coordinate on Δ denote as z_1 .

Set

$$a_f^\epsilon(z_1) = \frac{i}{2} \int_{\Delta_{z_1}(r_1)} t_{f,\epsilon}^{n\bar{n}} dz_n \wedge d\bar{z}_n,$$

where $t_{f,\epsilon}^{n\bar{n}}$ is the smoothing of $t_f^{n\bar{n}}$ by convolution. Since $t_{f,\epsilon}^{n\bar{n}} \rightarrow t_f^{n\bar{n}}$ in L^1_{loc} we get by Fubini Theorem that $a_f^\epsilon \rightarrow a_f$ in L^1_{loc} . Therefore using (7.3) for dimension two we obtain

$$\begin{aligned} \langle \Delta a_f^\epsilon, \varphi \rangle &= \frac{i}{2} \int_{\Delta(\delta, z_j)} \varphi(z_1) \left(\int_{\Delta_{z_1}(r_1)} \frac{\partial^2 t_{f,\epsilon}^{n\bar{n}}}{\partial z_1 \partial \bar{z}_1} dz_n \wedge d\bar{z}_n \right) dz_1 \wedge d\bar{z}_1 = \\ &= \frac{i}{2} \int_{\mathbb{D}(\delta, z_j)} \varphi(z_1) \left(\frac{i}{2} \int_{\partial \Delta_{z_1}(r_1)} \frac{\partial t_{f,\epsilon}^{1\bar{2}}}{\partial \bar{z}_1} d\bar{z}_2 + \frac{i}{2} \int_{\partial \mathbb{D}_{z_1}(r_1)} \frac{\partial t_{f,\epsilon}^{2\bar{1}}}{\partial z_1} dz_2 \right) dz_1 \wedge d\bar{z}_1 - \\ &\quad - \frac{i}{2} \int_{\mathbb{D}(\delta, z_j)} \varphi(z_1) \left(\int_{\partial \mathbb{D}_{z_1}(r_1)} \frac{\partial t_{f,\epsilon}^{1\bar{1}}}{\partial \bar{z}_2} d\bar{z}_2 \right) dz_1 \wedge d\bar{z}_1 \longrightarrow \langle \phi_f, \varphi \rangle \end{aligned}$$

as $\epsilon \rightarrow 0$. Therefore, $\Delta a_f = \phi_f$ in Δ in the sense of distributions. By regularity of the Laplacian $a_f \in C^\infty$ on Δ and therefore $c_A(f) = 0$ for all A and all f . Therefore a_f are smooth on Δ^{n-1} and $a_f = \psi_f + h_f$ there. ψ_f -s are equicontinuous and h_f are pluriharmonic everywhere and uniformly bounded from below. Moreover a_f are equicontinuous on Δ_ϵ^{n-1} by assumption. Therefore h_f are equicontinuous on Δ_ϵ^{n-1} . This implies equicontinuity of h_f on compacts of Δ^{-1} , and therefore the equicontinuity of a_f . Step 2 and therefore our Lemma are proved. \square

Lemmas 7.1 and 7.2 obviously imply the Theorem 5 from Introduction.

8. FATOU COMPONENTS

8.1. Case of dimension two and Fatou sets. Let us first prove the following

Proposition 8.1. *Suppose that a weakly converging sequence $\{f_k\}$ of meromorphic mappings from a two-dimensional domain U to a compact complex surface X doesn't converge strongly. Then X is bimeromorphic to \mathbb{P}^2 .*

Proof. Indeed, in that case there exists a point $a \in U$ and a neighborhood $V \ni a$ such that f_k converge uniformly on compacts of $V \setminus \{a\}$ but Γ_{f_k} do not converge to Γ_f , where $f : U \rightarrow X$ is the limit map. $\text{Vol}(\Gamma_{f_k})$ are uniformly bounded. Indeed for Kähler X it was proved in [Iv2] using Skoda's potential. In [Ne] it was proved for non-Kähler X using that fact that such X can contain only finitely many rational curves as well as existence of certain dd^c -exact $(2,2)$ -forms.

Therefore we see that the limit $\lim \Gamma_{f_k}$ contains Γ_f plus $\{a\} \times X$ (with some multiplicity). But this is a bubble and therefore X is rationally connected by Theorem 5. From the classification of surfaces we know that such X must be bimeromorphic to \mathbb{P}^2 . \square

Now let us prove Corollary 2 from Introduction. We start with the following lemma. For a meromorphic map $f : U \rightarrow X$ denote by

$$D_f := \pi_1(\{(z, x) \in U \times X : \dim_{(z, x)} \pi_2^{-1}(x) \geq 1\})$$

the set of degeneration of f . Here $\pi_1 : U \times X \rightarrow U$ and $\pi_2 : U \times X \rightarrow X$ are canonical projections. $f : U \rightarrow X$ is degenerate if $D_f = U$.

Lemma 8.1. *Let $f : X \rightarrow X$ be a nondegenerate meromorphic self-map of a compact complex surface X , let $z \in \Phi_s$ (resp. Φ_w). Then for every $l \geq 1$ one has*

$$f^l[z] \setminus f^l|_{D_{f^l}}(D_{f^l}) \subset \Phi_s \quad (\text{resp. } \Phi_w). \quad (8.1)$$

Proof. Let $a \in f^l[z] \setminus f^l|_{D_{f^l}}(D_{f^l})$, then there exist neighborhoods $U \ni z$ and $V \ni a$ such that $\pi_2 : (V \times U) \cap \Gamma_{f^l} \rightarrow U$ is proper. That means that $(f^l)^{-1} : U \rightarrow V$ is well defined as multivalued holomorphic map. Now let $\{f^{n_k}\} \subset \{f^n\}$ be a subsequence. By assumption from the sequence $\{f^{n_k+l}\}$ we can subtract a strongly/weakly converging on U subsequence $\{f^{n_{k_j}+l}\}$. That means that $\{f^{n_{k_j}} = f^{n_{k_j}+l} \circ f^l\}$ will converge in an appropriate sense on V . \square

Since every compact complex surface admits a dd^c -closed metric form Theorem 3 applies in our case and gives local pseudoconvexity of the weak Fatou set Φ_w . Suppose now that $\Phi_s \neq \Phi_w$.

a) By Proposition 8.1 $X = \mathbb{P}^2$.

b) There exists a point $p \in X$, a ball \mathbb{B} centered at p , a subsequence of iterates $\{f^{n_k}\}$, which uniformly converge on compacts of $\bar{B} \setminus \{p\}$ to a meromorphic map $f_\infty : \mathbb{B} \rightarrow X$, holomorphic on $B \setminus \{p\}$, but not converge strongly on any neighborhood of p . In particular this means that $p \in I(f_\infty)$ by Rouché Principle of [Iv2] and, moreover, $C = f_\infty[p]$ is a chain of rational curves $\bigcup_{i=1}^N C_i$. It was proved in [Iv2] for Kähler X and in [Ne] for non-Kähler case that $\text{Vol}(\Gamma_{f^{n_k}})$ are uniformly bounded on \bar{B} . So they are converging (after going to subsequence) in cycle topology to $\Gamma_{f_\infty} \cup d(\{p\} \times X)$ for some integer $d \geq 1$. Take a point $q \in X \setminus C$. Then for $k \gg 1$ $q \in f^{n_k}(\mathbb{B} \setminus \{p\})$. If moreover $q \notin f^{n_k}(D(f^{n_k}))$ then $q \in \Phi_w$. But $\bigcup_k f^{n_k}(D(f^{n_k}))$ is at most countable set of points and Φ_w is Levi-pseudoconvex. So $\Phi_w \supset X \setminus C$. Again from pseudoconvexity of Φ_w it follows that if Φ_w intersects some irreducible component of C then it contains this component minus the rest of C . I.e., $\Phi_w = \mathbb{P} \setminus \{\text{some components of } C\}$.

c) Take a point $(p, x) \in \{p\} \times X$ such that $x \in C$. Suppose that $\Gamma_{f_\infty} \cap (X \times \{x\})$ has (p, x) as isolated point. Then we can find a neighborhoods $W \ni p$ and $V \ni x$ with

$(\bar{W} \times \partial V) \cap \Gamma_{f_\infty} = \emptyset$. Thus $(\bar{W} \times \partial V) \cap \Gamma_{f^{n_k}} = \emptyset$ for k big enough. Therefore $\Phi_w \ni V$ as before and, moreover, Φ_w contains the component of C passing through x minus the rest of C .

Let us distinguish two cases.

Case 1. *Every component of C contains such a point.* In this case our sequence $\{f^{n_k}\}$ strongly converges on $X \setminus \{p_1, \dots, p_N\}$. Furthermore $\text{Vol} \Gamma_{f^{n_k}}$ are uniformly bounded but not less than $N \cdot \text{Vol} X$. This is possible only if f has degree one, $N = 1$ and f_∞ is degenerate map onto C . C in this case should consist only from one component.

Case 2. There exists a component C_1 of C such that for all points $x \in C_1 \setminus \cup_{i \neq 1} C_i$ $\dim_{(p,x)} \Gamma_{f_\infty} \cap (X \times \{x\}) > 0$. f_∞ is a degenerate mapping of X onto C_1 and therefore again C_1 is a single component of C . I.e., in both cases C consists from one rational curve only. □

The following simple example shows that the situation described in part (ii) of this theorem can really happen. Let $X = \mathbb{CP}^2$ and $f : [z_0 : z_1 : z_2] \rightarrow [z_0 : 2z_1 : 2z_2]$. Then for this f we have the phenomena described above with $p = [1 : 0 : 0]$ and $C = \{z_0 = 0\}$.

8.2. Example. Another example relevant to the Fatou sets is the following

Example 8.1. Consider the following rational self-map \mathbb{P}^2 :

$$f : [z_0 : z_1 : z_2] \rightarrow [z_0^2 z_1 : z_1^3 : z_0^2 z_2]. \quad (8.2)$$

By induction one easily checks that

$$f^k : [z_0 : z_1 : z_2] \rightarrow [z_0^{2^k} z_1^{2^k-1} : z_1^{2^{k+1}-1} : z_0^{2^{k+1}-2} z_2]. \quad (8.3)$$




FIGURE 1. Mapping f contracts the line $l_1 := \{z_1 = 0\}$ to the first of its points of indeterminacy $q = [0 : 0 : 1]$, line at infinity $l_0 := \{z_0 = 0\}$ to the regular point $r = [0 : 1 : 0]$ and do not contracts anything to its second point of indeterminacy $p = [1 : 0 : 0]$. Levi flat cone = Julia set for f is marked by two punctured lines.

Cover \mathbb{P}^2 by three standard affine charts $U_i = \{z_i \neq 0\}$ with coordinates $u_1 = \frac{z_1}{z_0}, u_2 = \frac{z_2}{z_0}$, $v_1 = z_0/z_1, v_2 = z_2/z_1$ and $w_1 = z_0/z_2, w_2 = z_1/z_2$ respectively. Mapping $f : U_0 \rightarrow U_0$ writes as

$$f : (u_1, u_2) \rightarrow (u_1^2, u_2/u_1). \quad (8.4)$$

We see from here that f has degree 2. Furthermore f^k writes as

$$f^k : (u_1, u_2) \rightarrow (u_1^{2^k}, u_2/u_1^{2^k-1}). \quad (8.5)$$

In the charts $f : U_1 \rightarrow U_1$ our iterate writes as

$$f^k : (v_1, v_2) \rightarrow (v_1^{2^k}, v_1^{2^{k+1}-2}v_2) \rightarrow (0, 0) = r \quad \text{on} \quad \{|v_1| < 1\}. \quad (8.6)$$

I.e., we see that $\Phi^1 = \{|v_1| < 1\} = \{|u_1| > 1\}$ is a component of the Fatou set of f , in all senses, because all f^k are holomorphic there and converge uniformly on compacts to a constant map to $r = [0 : 1 : 0]$.

Levi flat cone $L := \{|z_0| = |z_1|\}$ is a Julia set of f . It contains one of two indeterminacy points of f , namely $q = [0 : 0 : 1]$. A connected component Ω of $\mathbb{P}^2 \setminus L$ different from Φ carries a more interesting information about f^k . First of all remark that as a mapping from U_2 to U_2 our iterate writes as

$$f^k : (w_1, w_2) \rightarrow \left(\frac{w_2^{2^k-1}}{w_1^{2^k-2}}, \frac{w_2^{2^{k+1}-1}}{w_1^{2^{k+1}-2}} \right) \rightarrow (0, 0) = q \quad \text{on} \quad \{|w_2| < |w_1|\} = \{|u_1| < 1, u_2 \neq 0\}. \quad (8.7)$$

Therefore the second component Φ_s^2 of the strong Fatou set contains the domain $\Omega \setminus \{u_2 = 0\}$. Since it is easy from (8.5) to see that f^k on compacts in $\Omega \setminus \{u_2 = 0\}$ converge to q , and on the puncture disc $\Delta^* := \{u_2 = 0, 0 < |u_1| < 1\}$ to p , we conclude that $\Phi_s^2 = \Phi_w^2 = \Omega \setminus \{u_2 = 0\}$. Remark that the second component Φ^2 of f in the sense of [FS] is smaller, namely it is equal to $\Omega \setminus (\{u_2 = 0\} \cup \{u_1 = 0\})$, because the projective line $l_1 := \{z_1 = 0\}$ is the preimage of I_f (and of all I_{f^k}) under f . Now let us turn to the second component Φ_Γ^2 of the Γ -Fatou set of f .

Lemma 8.2. *For a fixed $0 < \varepsilon < 1$ the volumes of graphs of f_k over the bidisk $\Delta_\varepsilon^2 \subset U_0$ centered at p are uniformly bounded. In particular $\Phi_\Gamma^2 = \Omega$.*

Proof. To estimate the volume of Γ_{f_k} over a neighborhood of p we use coordinates u_1, u_2 and representation (8.5). In these coordinates $\Delta_\varepsilon^2 = \{u : \|u\| < \varepsilon\}$. Since f^k preserves the vertical lines $\{u_1 = \text{const}\}$ we can simplify our computations assuming that f takes values in $\Delta \times \mathbb{P}^1$, the last being equipped with the Hermitian metric form $\omega = \omega_1 + \omega_2 = \frac{i}{2} dz_1 \wedge d\bar{z}_1 + \frac{i}{2} \frac{dz_2 \wedge d\bar{z}_2}{(1+|z_2|^2)^2}$. Now we get

$$\begin{aligned} (f^k)^* \omega &= \frac{i}{2} \left[2^{2k} |u_1|^{2^{k+1}-2} + \frac{(1-2^k)^2 |u_1|^{2^{k+1}-4}}{(|u_1|^{2^{k+1}-2} + |u_2|^2)^2} \right] du_1 \wedge d\bar{u}_1 + \frac{i}{2} |u_1|^{2^{k+1}-2} \frac{du_2 \wedge d\bar{u}_2}{(|u_1|^{2^{k+1}-2} + |u_2|^2)^2} \\ &+ \frac{i}{2} (1-2^k) \bar{u}_1^{-2k} |u_1|^{2^{k+2}-4} \frac{du_2 \wedge d\bar{u}_1}{(|u_1|^{2^{k+1}-2} + |u_2|^2)^2} + \frac{i}{2} (1-2^k) u_1^{-2k} |u_1|^{2^{k+2}-4} \frac{du_1 \wedge d\bar{u}_2}{(|u_1|^{2^{k+1}-2} + |u_2|^2)^2}. \end{aligned}$$

Therefore

$$\begin{aligned} &\int_{\Delta_\varepsilon^2} (f^k)^* \omega \wedge dd^c \|u\|^2 = \\ &= 4\pi^2 \int_0^\varepsilon \int_0^\varepsilon \left[2^{2k} r_1^{2^{k+1}-1} + (1-2^k)^2 \frac{r_1^{2^{k+1}-3}}{(r_1^{2^{k+1}-2} + r_2^2)^2} + \frac{r_1^{2^{k+1}-1}}{(r_1^{2^{k+1}-2} + r_2^2)^2} \right] dr_1 r_2 dr_2 \leq \\ &\leq \frac{4\pi^2 \varepsilon^{2^{k+1}}}{2^{2^{k+1}-2k}} + 2\pi^2 2^{2k} \int_0^\varepsilon \int_0^{\varepsilon^2} \frac{r_1^{2^{k+1}-3} t}{(r_1^{2^{k+1}-2} + t)^2} dt dr_1 + 2\pi^2 2^{2k} \int_0^\varepsilon \int_0^{\varepsilon^2} \frac{r_1^{2^{k+1}-1}}{(r_1^{2^{k+1}-2} + t)^2} dt dr_1 \leq \end{aligned}$$

$$\begin{aligned}
&\leq 2\pi^2 2^{2k} \int_0^\varepsilon \int_0^{\varepsilon^2} \frac{r_1^{2^{k+1}-3}}{r_1^{2^{k+1}-2} + t} dt dr_1 + 2\pi^2 \int_0^\varepsilon \left(\frac{-1}{r_1^{2^{k+1}-2} + t} \Big|_0^{\varepsilon^2} r_1^{2^{k+1}-1} \right) dr_1 \leq \\
&\leq \pi^2 2^{3k+1} \int_0^\varepsilon r_1^{2^{k+1}-3} \ln \frac{1}{r_1} dr_1 + 2\pi^2 \int_0^\varepsilon r_1 dr_1.
\end{aligned}$$

The second term is bounded and doesn't tend to zero as $k \rightarrow +\infty$. The first for $0 < \varepsilon < 1$ can be obviously bounded by

$$\pi^2 2^{3k+1} \int_0^\varepsilon r^{2^{k+1}-4} dr = \frac{\pi^2 2^{3k+1}}{2^{k+1}-3} \varepsilon^{2^{k+1}-3} \rightarrow 0$$

as $k \rightarrow +\infty$. And finally

$$\begin{aligned}
&\int_{\Delta_\varepsilon^2} (f^k)^* \omega = \int_{\Delta_\varepsilon^2} \frac{2^{2k} |u_1|^{2^{k+2}-4}}{(|u_1|^{2^{k+1}-2} + |u_2|^2)^2} + \frac{(2^k - 1)^2 |u_1|^{2^{k+2}-6} |u_2|^2}{(|u_1|^{2^{k+1}-2} + |u_2|^2)^4} + \\
&+ \frac{2(2^k - 1)^2 |u_1|^{2^{k+3}-8} |u_2|^2}{|u_1|^{2^{k+1}} (|u_1|^{2^{k+1}-2} + |u_2|^2)^4} d^4 m(u) \leq \int_{\Delta_\varepsilon^2} \frac{2^{2k+2} |u_1|^{2^{k+2}-6}}{(|u_1|^{2^{k+1}-2} + |u_2|^2)^2} d^4 m(u) \approx \\
&\approx 2^{2k+2} \int_0^\varepsilon \int_0^{\varepsilon^2} \frac{r_1^{2^{k+2}-5}}{(r_1^{2^{k+1}-2} + t)^2} dt dr_1 \leq 2^{2k+2} \int_0^\varepsilon r^{2^{k+1}-3} dr = \frac{2^{2k+2}}{2^{k+1}-2} \varepsilon^{2^{k+1}-2} \rightarrow 0
\end{aligned}$$

as $k \rightarrow +\infty$. Therefore the lemma is proved. \square

Remark that the integral of $(f^k)^* \omega^2$ degenerates, as it should be, because f^k Γ -converges on Ω to a constant map. And to the contrary the integral of $(f^k)^* \omega$ doesn't degenerate, moreover has order ε^2 . That means that bubbling takes place over all points of the disk $\Delta^* = \{u_2 = 0, 0 < |u_1| < 1\}$. We see that for our map one has

$$\Phi \subset \Phi_w = \Phi_s \subset \Phi_\Gamma,$$

and inclusions are strict.

Finally let us see what is going on over the indeterminacy point $p = (0, 0)$ in coordinates (u_1, u_2) . Blowing up $2^k - 1$ times at zero we see that f^k stays degenerate on all exceptional curves except the last one, which it send onto $l_1 = \{u_1 = 0\}$. In appropriate coordinates v_1, v_2 on the last blow up f^k writes as

$$\begin{cases} u_1 = v_1^{2^k} \\ u_2 = v_2. \end{cases}$$

Therefore the dynamical picture over p can be described as follows. Let $\hat{\Omega}$ be an infinite blow up of Ω over p and let $C = \bigcup_{i=1}^\infty C_i$ be the Nori string of rational curves on $\hat{\Omega}$ over p . Then every f^k lifts to a holomorphic map $\hat{f}^k : \hat{\Omega} \rightarrow \mathbb{P}^2$ which is constantly equal to q on every C_i except C_{2^k-1} . The last curve it sends bijectively onto the line l_1 . At the same time in the sense of divisors (currents) $f^k(C_{2^k-1}) = 2^k l_1$.

Remark 8.1. a) One can in the same way produce mappings of any given degree with the same properties as in Example 8.1. It is sufficient to take

$$f : [z_0 : z_1 : z_2] \rightarrow [z_0^d z_1 : z_1^{d+1} : z_0^d z_2]. \quad (8.8)$$

b) Let us quote the result of Maegava, see [M], which shows that under an additional assumption of "algebraic stability" of the dominant rational self-map f the Fatou set of [FS] coincides with Φ_s and Φ_w .

REFERENCES

- [Ar] ARAUJO C.: *Rationally connected varieties*. Contemp. Math. **388**, 1-16 (2006).
- [CT] CARTAN B., THULLEN P.: *Zur der Singularitäten der Funktionen mehrerer Verandenlichen: Regularitäts und Konvergenzbereiche*. Math. Ann. **106**, 617-647 (1932).
- [Ba] BARLET D.: *Espace analytique reduit des cycles analytiques complexes compacts d'un espace analytique complexe de dimension finie*. Seminar Norguet IX, Lect. Notes Math., **482**, 1-157, (1975).
- [BT] BEDFORD E., TAYLOR B.A.: *A new capacity for plurisubharmonic functions*. Acta Math. **149**, no. 1-2, 1-40 (1982).
- [DG] DOCQUIER F., GRAUERT H.: *Levisches Problem und Rungescher Satz für Teilgebiete Steinscher Mannigfaltigkeiten*. Math. Ann. **140**, 94-123, (1960).
- [FS] FORNAESS J.-E., SIBONY N.: *Complex dynamics in higher dimension. II*. Modern methods in complex analysis (Princeton, NJ, 1992), 135182, Ann. of Math. Stud., **137**, Princeton Univ. Press, Princeton, NJ, (1995).
- [Fj] FUJIKI A.: *Closedness of the Douady Space of Compact Kähler Spaces*. Publ. RIMS, Kyoto Univ. **14**, 1-52, (1978).
- [Fu] FUJIMOTO H.: *On families of meromorphic maps into the complex projective space*. Nagoya Math. J. **54**, 21-51, (1974).
- [Ga] GAUDUCHON P.: *Les metriques standard d'une surface a premier nombre de Betti pair*. Asterisque. Soc. Math. France. **126**, 129-135, (1985).
- [GK] GRIFFITHS P., KING J.: *Nevanlinna theory and holomorphic mappings between algebraic varieties*. Acta Math. **130**, 145-220, (1973).
- [Gr] GREEN M.: *The hyperbolicity of the complement of $2n+1$ hyperplanes in general position in \mathbb{P}^n , and related results*. Proc. Amer. Math. Soc. **66**, 109-113 (1977).
- [HS] HARVEY R., SHIFFMAN B.: *A Characterization of Holomorphic Chains*. Annals of Math. **99**, no: 3, 553-587 (1974).
- [Ho] HÖRMANDER L.: *An introduction to complex analysis in several variables*. Third edition, North-Holland Mathematical Library, **7**, North-Holland Publishing Co., Amsterdam, (1990).
- [Iv1] IVASHKOVICH S.: *The Hartogs-type extension theorem for the meromorphic maps into compact Kähler manifolds*. Invent. math. **109**, 47-54, (1992).
- [Iv2] IVASHKOVICH S.: *On convergence properties of meromorphic function and mappings (in russian)*. In "Complex Analysis in Modern Mathematics", B. Shabat memorial volume, FAZIS, Moscow, 133-151, (2001).
- [Iv3] IVASHKOVICH S.: *Extension properties of meromorphic mappings with values in non-Kähler complex manifolds*. Annals of Math. **160**, 795-837, (2004).
- [Iv4] IVASHKOVICH S.: *On fixed points of rational self-maps of mplex projective plane*. To appear in Indiana Univ. Math. J., see also arXiv:0911.5084 .
- [IS1] IVASHKOVICH S.; SHEVCHISHIN V.: *Structure of the moduli space in a neighborhood of a cusp-curve and meromorphic hulls*. Sbornik Math. **189** no. 9-10, 1335-1359 (1998).
- [IS2] IVASHKOVICH S.; SHEVCHISHIN V.: *Deformations of non-compact complex curves and envelopes of meromorphy of spheres*. Invent. Math. **136**, 571-602, (1999).

- [IS3] IVASHKOVICH S.; SHEVCHISHIN V.: *Gromov Compactness Theorem for J-Complex Curves with Boundary*. Int. Math. Res. Not. **22** 1167-1206, (2000).
- [Kg] KING J.: *A residue formula for complex subvarieties*. Proc. Carolina Conf. on Holomorphic Mappings and Minimal Surfaces (Chapel Hill, N.C., 1970) pp. 4356 Dept. of Math., Univ. of North Carolina.
- [Ki] KIERNAN P.: *Hyperbolically Imbedded Spaces and the Big Picard Theorem*. Math. Ann. **204**, 203-209, (1973).
- [Ks] KISELMAN C.: *Sur la définition de l'opérateur de Monge-Ampère complexe*. Lecture Notes in Math., Springer, Berlin, **1094**, 139-150, (1984).
- [Ko] KOBAYASHI S.: *Intrinsic distances, measures and geometric function theory*. BAMS **82**, no: 3, 357-416, (1976).
- [M] MAEGAWA K.: *Ascoli-Arzelà type theorem for rational iterations of the complex projective plane*. Proc. AMS **136**, no: 8, 2875-2879 (2008).
- [Me] MENEGHINI C.: *Renormalizing iterated repelling germs of \mathbb{C}^2* . Mat. Vesnik **56**, no. 3-4, 85-90, (2004).
- [Ne] NEJI F.: *Equicontinuous families of meromorphic mappings with values in compact complex surfaces*. To appear in Comp. Var. and Ell. Eq., see math.CV/1102.3608 .
- [Ni] NISHINO T.: *Sur les familles de surfaces analytiques*. J. Math. Kyoto Univ. **1** 357-377, (1962).
- [NW] NOGUCHI J., WINKELMAN J.: *Holomorphic curves and integral points of divisors*. Math. Z. **239** 593-610, (2002).
- [R] RANSFORD T.: *Potential Theory in the Complex Plane*. London Mathematical Society Student Texts 28, Cambridge Univ. Press, Cambridge, UK, (1995).
- [Rm] RIEMENSCHNEIDER O.: *ber den Flächeninhalt analytischer Mengen und die Erzeugung kpseudokonvexer Gebiete*. Invent. Math. **2**, 307331, (1967).
- [Ru] RUTISHAUSER H.: *Über die Folgen und Scharen von analytischen und meromorphen Funktionen mehrerer variablen, sowie von analytischen und meromorphen abbildungen*. Acta Math. **83** 249-325, (1950).
- [Si] SIU Y.-T.: *Extension of meromorphic maps into Kähler manifolds*. Annals of Math. **102**, 421-462, (1975).
- [Sk] SKODA H.: *Sous-ensembles analytiques d'ordre fini ou infini dans \mathbb{C}^n* . Bulletin de la S.M.F., **100**, 353-408, (1972).
- [Stl] STOLL W. *Normal families of non-negative divisors*. Math. Z., **84**, 154-218, (1964).
- [St] STOLZENBERG G.: *Volumes, Limits and Extension of Analytic Varieties* Springer Verlag (1966).
- [Za] ZAIDENBERG M.: *Picard's theorem and hyperbolicity*. Siberian. Math. J. **24**, no:6, 858-867, (1983).

UNIVERSITÉ DE LILLE-1, UFR DE MATHÉMATIQUES, 59655 VILLENEUVE D'ASCQ, FRANCE
E-mail address: ivachkov@math.univ-lille1.fr

UNIVERSITÉ DE LILLE-1, UFR DE MATHÉMATIQUES, 59655 VILLENEUVE D'ASCQ, FRANCE
E-mail address: Fethi.neji@math.univ-lille1.fr